

Non-homogeneous locally free actions of the affine group

Masayuki ASAOKA *

Department of Mathematics, Kyoto University

April 24, 2009

Abstract: We classify locally free smooth actions of the real affine group on closed three-dimensional manifolds up to smooth conjugacy. In particular, we show that there are non-homogeneous actions when the manifold is not a rational homology sphere and its fundamental group is not solvable.

Keywords: locally free actions of Lie groups, Anosov foliations.

1 Introduction

Let GA be the group of orientation preserving affine transformations of the real line. The main theme of this paper is a classification of locally free actions of GA on closed three-dimensional manifolds. In 1979, Ghys [9] showed that any volume-preserving action is smoothly conjugate to a homogeneous one. Existence or non-existence of non-homogeneous actions was open for thirty years. In this paper, we give a complete classification of actions up to smooth conjugacy. As a corollary, there *exists* a non-homogeneous one.

Let G be a Lie group and M a C^∞ manifold. We say a C^∞ right action $\rho : M \times G \rightarrow M$ is *locally free* if the isotropy subgroup $\{g \in G \mid \rho(p, g) = p\}$ is discrete in G for any $p \in M$. By $\mathcal{O}_\rho(p)$, we denote the orbit $\{\rho(p, g) \mid g \in G\}$ of $p \in M$. When the action ρ is locally free, the decomposition $\mathcal{O}_\rho = \{\mathcal{O}_\rho(p) \mid p \in M\}$ of M is a C^∞ foliation.

We say two right actions ρ_1 and ρ_2 of a Lie group G on manifolds M_1 and M_2 are C^∞ *conjugate* if there exist an isomorphism σ of G and a C^∞ diffeomorphism H from M_1 to M_2 such that

$$H(\rho_1(p, g)) = \rho_2(H(p), \sigma(g))$$

for any $g \in G$ and $p \in M_1$. The map H is called a C^∞ *conjugacy* between ρ_1 and ρ_2 . We also say ρ_1 and ρ_2 are C^∞ *orbit equivalent* if there exists a C^∞

*Partially supported by JSPS PostDoctoral Fellowships for Research Abroad and Grant-in-Aid for Encouragement of Young Scientists (B)

diffeomorphism H from M_1 to M_2 such that

$$H(\mathcal{O}_{\rho_1}(p)) = \mathcal{O}_{\rho_2}(H(p))$$

for any $p \in M_1$.

There are two classical examples of locally free actions of GA on closed three-dimensional manifolds. Let $\widetilde{\text{PSL}}(2, \mathbb{R})$ be the universal covering group of the group $\text{PSL}(2, \mathbb{R})$ of projective transformations of the real projective line $\mathbb{RP}^1$. It contains a closed subgroup H which is isomorphic to GA, *e.g.*, the group of lower triangle elements. Fix an isomorphism $i : \text{GA} \rightarrow H$. For each cocompact lattice Γ , we define a locally free action ρ_1 of GA on $\Gamma \backslash \widetilde{\text{PSL}}(2, \mathbb{R})$ by $\rho_1(\Gamma g, h) = \Gamma(g \cdot i(h))$. The same construction gives another locally free action ρ_2 when we start with a three-dimensional solvable subgroup

$$\text{Solv} = \{h_{t,x,y} : (u, v) \mapsto (e^{-t}u + x, e^t v + y) \mid t, x, y \in \mathbb{R}\}$$

of the group of affine transformations of $\mathbb{R}^2$. In this case, GA can be identified with the subgroup $\{h_{t,0,y} \mid t, x \in \mathbb{R}\}$. We call an action of GA *homogeneous* if it is C^∞ conjugate to one of the above actions.

As mentioned at the beginning, Ghys showed the rigidity of volume-preserving actions in his doctoral thesis [9].

Theorem 1.1 ([9]). *If a C^∞ locally free action of GA on a closed and connected three-dimensional manifold admits a continuous invariant volume, then it is homogeneous.*

The following problem is quite natural, but was open for thirty years.

Problem 1.2. Remove the assumption on an invariant volume, or construct a C^∞ locally free action of GA on a closed three-dimensional manifold which is not homogeneous.

See [4, p.2]. A related problem was also mentioned in [23, p.1841].

There are several generalizations of Theorem 1.1 to codimension-one volume-preserving actions of higher-dimensional solvable groups ([3], [4], [6], [27], and [28]). Recently, Belliard [5] announced that the assumption on an invariant volume is unnecessary for a large class of higher-dimensional solvable groups. However, his result does not cover actions of GA.

The group GA is generated by one-parameter subgroups $\{X^t\}_{t \in \mathbb{R}}$ and $\{U^y\}_{y \in \mathbb{R}}$ of GA which satisfy

$$U^y X^t = X^t U^{(\exp t) \cdot y}. \quad (1)$$

For a right action ρ of GA on a manifold M , let $\Phi_\rho = \{\Phi_\rho^t\}_{t \in \mathbb{R}}$ be the flow defined by the ρ -action of $\{X^t\}_{t \in \mathbb{R}}$ on M , *i.e.*, $\Phi_\rho^t(p) = \rho(p, X^t)$. Ghys showed that the Jacobian $\det D\Phi_\rho^t$ is controlled by a closed one-form.

Theorem 1.3 ([11, Chapter IV]). *Let M be a three-dimensional closed Riemannian manifold and $\rho : M \times \text{GA} \rightarrow M$ a C^∞ locally free action. Then, there exists a closed one-form ω on M such that*

$$(\Phi_\rho^t)^* \omega - \omega = d(\log \det D\Phi_\rho^t). \quad (2)$$

for any $t \in \mathbb{R}$.

As a corollary, he obtained a partial answer to Problem 1.2.

Corollary 1.4 ([11]). *Any C^∞ locally free action of GA on a rational homology sphere preserves a continuous volume. In particular, it is a homogeneous action.*

The author of this paper also showed the rigidity of the orbit foliation.

Theorem 1.5 ([1]). *Let ρ be a C^∞ locally free action of GA on a three-dimensional, closed, and connected manifold M . Then, Φ_ρ is an Anosov flow and ρ is C^∞ orbit equivalent to a homogeneous action. In particular, $M = \Gamma \backslash \mathrm{PSL}(2, \mathbb{R})$ or $\Gamma \backslash \mathrm{Solv}$ with a cocompact lattice Γ .*

The above results and the classification of Anosov flows on solvable manifolds due to Plante [24] give the answer to Problem 1.2 for the case that the fundamental group of M is solvable.

Theorem 1.6 (Ghys). *Let Γ be a cocompact lattice of Solv . Then, any C^∞ locally free action of GA on $\Gamma \backslash \mathrm{Solv}$ preserves a continuous volume. In particular, it is a homogeneous action.*

We will give the proof in Appendix since it has not been published.

The only unsolved case is that the underlying manifold is $\Gamma \backslash \mathrm{PSL}(2, \mathbb{R})$ with a cocompact lattice Γ and it is not a rational homology sphere. The main goal of this paper is to give a complete classification of C^∞ locally free actions of GA in this case. As a consequence, we will see the existence of non-homogeneous actions.

For a cocompact lattice Γ of $\mathrm{PSL}(2, \mathbb{R})$, we denote the quotient space $\Gamma \backslash \mathrm{PSL}(2, \mathbb{R})$ by M_Γ . It admits a natural homogeneous action ρ_Γ (see Section 3.1). Let $\mathcal{A}_\Gamma$ be the set of C^∞ locally free actions ρ of GA on M_Γ such that $\mathcal{O}_\rho = \mathcal{O}_{\rho_\Gamma}$. As Theorem 1.5, any C^∞ locally free action of GA on a three-dimensional, closed, and connected manifold with the non-solvable fundamental group is C^∞ conjugate to an action in $\mathcal{A}_\Gamma$ for some cocompact lattice Γ of $\mathrm{PSL}(2, \mathbb{R})$. Therefore, it is sufficient to classify actions in $\mathcal{A}_\Gamma$ up to C^∞ conjugacy.

As we will show in Proposition 3.7, the one-form ω in Theorem 1.3 is uniquely determined for each action in $\mathcal{A}_\Gamma$. Hence, the equation (2) defines a map $\bar{a}$ from $\mathcal{A}_\Gamma$ to the first de Rham cohomology group $H^1(M_\Gamma, \mathbb{R})$. The first theorem asserts that the map $\bar{a}$ parametrizes C^∞ conjugacy classes of $\mathcal{A}_\Gamma$.

Theorem A. *Two actions ρ_1 and ρ_2 in $\mathcal{A}_\Gamma$ is C^∞ conjugate via a diffeomorphism isotopic to the identity if and only if $\bar{a}(\rho_1) = \bar{a}(\rho_2)$.*

The second theorem describes the image of $\bar{a}$. Recall that the hyperbolic plane $\mathbb{H}^2$ admits a natural isometric action of $\mathrm{PSL}(2, \mathbb{R})$. We define the *translation length* $L(P)$ of $P \in \mathrm{PSL}(2, \mathbb{R})$ by

$$L(P) = \inf_{z \in \mathbb{H}^2} d_{\mathbb{H}^2}(z, Pz),$$

where $d_{\mathbb{H}^2}$ is the distance induced by the hyperbolic metric. We identify the lattice Γ with the fundamental group of M_Γ . Then, the evaluation $a(\gamma)$ is well-defined for a cohomology class a in $H^1(M_\Gamma, \mathbb{R})$ and $\gamma \in \Gamma$. For a cocompact lattice Γ of $\widetilde{\mathrm{PSL}}(2, \mathbb{R})$, let Δ_Γ be the set of cohomology classes a in $H^1(M_\Gamma, \mathbb{R})$ that satisfy

$$\sup \left\{ \left| \frac{a(\gamma)}{L(\gamma)} \right| \mid \gamma \in \Gamma, L(\gamma) \neq 0 \right\} < 1. \quad (3)$$

As we will see in Corollary 3.12, it is a non-empty, open, and convex subset of $H^1(M_\Gamma, \mathbb{R})$.

Theorem B.

$$\Delta_\Gamma = \{\bar{a}(\rho) \mid \rho \in \mathcal{A}_\Gamma\}$$

for any cocompact lattice Γ of $\widetilde{\mathrm{PSL}}(2, \mathbb{R})$. The following theorem completes the classification of actions in $\mathcal{A}_\Gamma$ up to C^∞ conjugacy.

The third theorem is concerned with real analytic realization of non-homogeneous actions.

Theorem C. Any action in $\mathcal{A}_\Gamma$ is C^∞ conjugate to a C^ω action.

Corollary. If M_Γ is not a rational homology sphere, then it admits a C^ω locally free action of GA which is not homogeneous.

Recall that the flow Φ_ρ is Anosov by Theorem 1.5. The last theorem shows that the smoothness of the weak stable foliation of Φ_ρ characterizes homogeneous actions.

Theorem D. Let Γ be a cocompact lattice of $\widetilde{\mathrm{PSL}}(2, \mathbb{R})$ and ρ an action in $\mathcal{A}_\Gamma$. If the weak stable foliation of the flow Φ_ρ is of class C^2 , then ρ is a homogeneous action.

To end the introduction, we pose some problems. First, the above results give the complete list C^∞ locally free actions of GA on three-dimensional closed manifolds. However, existence of smooth deformation of an action is still open.

Problem 1.7. Construct a C^∞ (or C^ω) family $\{\rho_a\}_{a \in \Delta_\Gamma}$ in $\mathcal{A}_\Gamma$ satisfying $\bar{a}_\Gamma(\rho_a) = a$.

Second, our construction of non-homogeneous actions is not ‘constructive’ since it uses Margulis measure, whose existence relies on the Schauder-Tychonoff fixed point theorem. So, the following question may be important to understand non-homogeneous actions.

Problem 1.8. Let X and U be the vector fields which generates the homogeneous actions of $\{X^t\}_{t \in \mathbb{R}}$ and $\{U^y\}_{y \in \mathbb{R}}$ on M_Γ , respectively. Take $\rho \in \mathcal{A}_\Gamma$ and let f_1, g_1, f_2 , and g_2 be functions such that vector fields $f_1 \cdot X + g_1 \cdot U$ and $f_2 \cdot X + g_2 \cdot U$ generate $\{\rho(X^t)\}_{t \in \mathbb{R}}$ and $\{\rho(U^y)\}_{y \in \mathbb{R}}$, respectively. Describe the functions f_1, g_1, f_2 , and g_2 explicitly.

Acknowledgements The author wrote the first version of this paper when he stayed at Unité de Mathématiques Pures et Appliquées, École Normale Supérieure de Lyon. He would like to thank the members of UMPA, especially Professor Étienne Ghys for their warm hospitality. The author is also grateful to Professor Shigenori Matsumoto for many suggestions to improve the paper.

2 Preliminary results

In this section, we review some basic results which we need later.

2.1 Anosov flows

In this subsection, we recall the definition and basic properties of Anosov flows. Let Φ be a C^1 flow on a closed manifold M without stationary points. Its differential defines a flow $D\Phi$ on the tangent bundle TM of M . Let $T\Phi$ be the one-dimensional subbundle of TM which is tangent to the orbits of Φ . The flow Φ is called *Anosov* if there exists a continuous $D\Phi$ -invariant splitting $TM = T\Phi \oplus E^{ss} \oplus E^{uu}$ and two constants $\lambda > 0$ and $C > 0$ such that

$$\max \{ \|D\Phi^t|_{E^{ss}(p)}\|, \|D\Phi^{-t}|_{E^{uu}(p)}\| \} < C \exp(-\lambda t)$$

for any $p \in M$ and $t \geq 0$, where $\|\cdot\|$ is the operator norm with respect to a Riemannian metric of M . The above splitting is unique for each Φ and is called *the Anosov splitting* of Φ . The $D\Phi$ -invariant subbundles E^{ss} , E^{uu} , $E^s = T\Phi \oplus E^{ss}$, and $E^u = T\Phi \oplus E^{uu}$ of Φ generate continuous foliations $\mathcal{F}^{ss}$, $\mathcal{F}^{uu}$, $\mathcal{F}^s$, and $\mathcal{F}^u$. They are called *the strong stable*, *strong unstable*, *weak stable*, and *weak unstable foliations*, respectively. By $\text{Per}(\Phi)$, we denote the union of periodic orbits of Φ . Let $\tau(p; \Phi)$ be the period of $p \in \text{Per}(\Phi)$ and put

$$\begin{aligned} J(p; \Phi) &= \log \det D\Phi^{\tau(p; \Phi)}(p), \\ J^s(p; \Phi) &= \log \det \left(D\Phi^{\tau(p; \Phi)}|_{E^{ss}(p)} \right), \\ J^u(p; \Phi) &= \log \det \left(D\Phi^{\tau(p; \Phi)}|_{E^{uu}(p)} \right). \end{aligned}$$

Remark that J , J^s and J^u does not depend on the choice of the metric $\|\cdot\|$.

Let Φ_i be a flow on a manifold M_i for $i = 1, 2$. We say a homeomorphism $H : M_1 \rightarrow M_2$ is a *topological conjugacy* if $H \circ \Phi_1^t = \Phi_2^t \circ H$ for any $t \in \mathbb{R}$. When H is a C^r diffeomorphism, it is called a *C^r conjugacy*. We say a homeomorphism $H : M_1 \rightarrow M_2$ is an *topological equivalence* if

$$H(\{\Phi_1^t(p) \mid t \geq 0\}) = \{\Phi_2^t(H(p)) \mid t \geq 0\}$$

for any $p \in M_1$. When H is a C^r diffeomorphism, it is called a *C^r equivalence*.

We say a function α on $M \times \mathbb{R}$ is a *cocycle* over a flow Ψ if $\alpha(p, t + t') = \alpha(\Psi^t(p), t') + \alpha(p, t)$. Recall that a flow is called *topologically transitive* if it

admits a dense orbit. The original Hölder continuous version of the following theorem is due to Livschitz, and the C^∞ version is due to de la Llave, Marco, and Moriyón.

The Livschitz Theorem ([19], [20]). *Let Φ be a C^2 topologically transitive Anosov flow on a closed manifold M and α be a Hölder continuous cocycle over Φ . If $\alpha(p, \tau(p; \Phi)) = 0$ for any $p \in \text{Per}(\Phi)$, then there exists a Hölder continuous function β such that*

$$\alpha(p, t) = \beta(p) - \beta \circ \Phi^t(p)$$

for any $p \in M$ and $t \in \mathbb{R}$. It is unique up to constant term. Moreover, if Φ and α are of class C^∞ , then also β is.

The above theorem has applications to the conjugacy problem of Anosov flows.

Theorem 2.1 ([21]). *Let Φ_1 and Φ_2 be C^∞ topologically transitive Anosov flows on three-dimensional closed manifolds M_1 and M_2 , respectively. Suppose that a topological conjugacy H between Φ_1 and Φ_2 satisfies $J^s(p; \Phi_1) = J^s(H(p); \Phi_2)$ and $J^u(p; \Phi_1) = J^u(H(p); \Phi_2)$ for any $p \in \text{Per}(\Phi_1)$. Then, H is a C^∞ conjugacy.*

Theorem 2.2 (c.f. [18, Theorem 19.2.9]). *Let Φ_1 and Φ_2 be C^∞ Anosov flows on closed manifolds M_1 and M_2 , respectively. Suppose that a topological equivalence H between Φ_1 and Φ_2 satisfies $\tau(p; \Phi_1) = \tau(H(p); \Phi_2)$ for any $p \in \text{Per}(\Phi_1)$. Then, there exists a continuous function β on M_1 such that a continuous map $H_\beta : M_1 \rightarrow M_2$ defined by $H_\beta(p) = \Phi_2^{\beta(p)}(H(p))$ is a topological conjugacy between Φ_1 and Φ_2 .*

Proof. As Theorem 19.1.5 of [18], we can replace a topological equivalence H by a bi-Hölder one. Then, there exists a Hölder continuous cocycle $\alpha : M_1 \times \mathbb{R} \rightarrow \mathbb{R}$ such that $H(\Phi_1^t(p)) = \Phi_2^{\alpha(p, t)}(H(p))$ for any $p \in M$ and $t \in \mathbb{R}$. The assumption implies $\alpha(p, \tau(p; \Phi_1)) = \tau(H(p); \Phi_2) = \tau(p; \Phi_1)$ for any $p \in \Phi_1$. By the Livschitz theorem, there exists a Hölder continuous function β such that $\alpha(p, t) = t + \beta(p) - \beta \circ \Phi_1(p)$. Since H is a topological equivalence, $\alpha(p, t) > 0$ if $t > 0$. It implies that H_β is locally injective. Since the mapping degree of H_β is one, it is a homeomorphism. Now, we have

$$\begin{aligned} H_\beta \circ \Phi_1^t(p) &= \Phi_2^{\beta \circ \Phi_1^t(p)} \circ H \circ \Phi_1^t(p) \\ &= \Phi_2^{\beta \circ \Phi_1^t(p) + \alpha(p, t)} \circ H(p) \\ &= \Phi_2^{t + \beta(p)} \circ H(p) = \Phi_2^t \circ H_\beta(p). \end{aligned}$$

□

The following is an inequality version of the Livschitz theorem.

Proposition 2.3. *Let Φ be a C^∞ Anosov flow on a closed manifold M , α be a C^∞ cocycle over Φ , and $\lambda > \lambda' > 0$ be constants. If $\alpha(p, \tau(p; \Phi)) \geq \lambda \cdot \tau(p; \Phi)$ for any $p \in \text{Per}(\Phi)$, then there exists a C^∞ function β such that*

$$\alpha(p, t) + \beta \circ \Phi^t(p) - \beta(p) \geq \lambda' t$$

for any $(p, t) \in M \times \mathbb{R}$.

Proof. First, we claim that there exist $T > 0$ such that $\alpha(p, T) \geq \lambda' T$ for any $p \in M$. Suppose that it does not hold. Then, there exist sequences $\{p_n \in M\}_{n \geq 1}$ and $\{t_n > 0\}_{n \geq 1}$ such that $\lim_{n \rightarrow \infty} t_n = \infty$ and $\alpha(p_n, t_n)/t_n < \lambda'$ for any $n \geq 1$. Put $f(p) = (\partial \alpha / \partial t)(p, 0)$. By taking a limit of the uniform measures on the orbit segment $\{\Phi^t(p) \mid t \in [0, t_n]\}$ and applying the ergodic decomposition theorem to it, we obtain an ergodic invariant probability measure μ of Φ such that

$$\int_M f(p) d\mu(p) \leq \liminf_{n \rightarrow \infty} \frac{1}{t_n} \int_0^{t_n} f \circ \Phi^t(p_n) dt = \liminf_{n \rightarrow \infty} \frac{\alpha(p_n, t_n)}{t_n} \leq \lambda'. \quad (4)$$

By a theorem due to Sigmund [26], any ergodic invariant probability measure of an Anosov flow can be approximated by invariant measures supported on periodic orbits. Hence, there exists a sequence $\{q_n \in \text{Per}(\Phi)\}_{n \geq 1}$ such that

$$\int_M f(p) d\mu(p) = \lim_{n \rightarrow \infty} \frac{1}{T_n} \int_0^{T_n} f \circ \Phi^t(q_n) dt = \lim_{n \rightarrow \infty} \frac{\alpha(q_n, T_n)}{\tau(q_n; T_n)} \geq \lambda, \quad (5)$$

where $T_n = \tau(q_n; \Phi)$. However, it contradicts (4). Therefore, the claim holds.

For $p \in M$, we define a function β by

$$\beta(p) = \frac{1}{T} \int_0^T \alpha(p, s) ds.$$

The identity

$$\alpha(p, s+t) = \alpha(\Phi^t(p), s) + \alpha(p, t) = \alpha(\Phi^s(p), t) + \alpha(p, s),$$

implies

$$\begin{aligned} \alpha(p, t) + \beta \circ \Phi^t(p) - \beta(p) &= \frac{1}{T} \int_0^T \alpha(\Phi^t(p), s) + \alpha(p, t) - \alpha(p, s) ds \\ &= \frac{1}{T} \int_0^T \alpha(\Phi^s(p), t) ds \\ &= \frac{1}{T} \int_0^T \left(\int_0^t f \circ \Phi^{s+s'}(p) ds' \right) ds \\ &= \int_0^t \frac{\alpha(\Phi^{s'}(p), T)}{T} ds' \geq \lambda' t. \end{aligned}$$

□

2.2 Topological entropy

Let Ψ be a continuous flow on a compact metric space (M, d) . For $t > 0$ and $\epsilon > 0$, we say a subset E of M is (t, ϵ) -separated if $\max_{0 \leq \tau \leq t} d(\Psi^\tau(p), \Psi^\tau(q)) \geq \epsilon$ for any mutually distinct elements $p, q \in E$. Let $S_\Psi(t, \epsilon)$ be the collection of all (t, ϵ) -separated subset of M . For a continuous cocycle α over Ψ , we put

$$N(\Psi, \alpha, t, \epsilon) = \sup_{E \in S_\Psi(t, \epsilon)} \left\{ \sum_{x \in E} \exp \alpha(x, t) \right\}.$$

We define the topological pressure $P(\alpha, \Psi)$ of a cocycle α with respect to Ψ by

$$P(f, \Psi) = \lim_{\epsilon \rightarrow 0} \limsup_{t \rightarrow \infty} \frac{1}{t} \log N(\Psi, \alpha, t, \epsilon).^1$$

The following lemma is easy to prove.

Lemma 2.4. *Suppose that two continuous cocycles α_1 and α_2 over a continuous flow Ψ , a continuous function β , and a real number C satisfies the identity*

$$\alpha_2(p, t) = \alpha_1(p, t) + \beta(p) - \beta \circ \Psi^t(p) + C \cdot t$$

Then, $P(\alpha_2, \Psi) = P(\alpha_1, \Psi) + C$. □

The topological entropy $h_{top}(\Psi)$ is defined by $h_{top}(\Psi) = P(0, \Psi)$. It is invariant under topological conjugacy, i.e., if two flows Ψ_1 and Ψ_2 are topologically conjugate, then $h_{top}(\Psi_1) = h_{top}(\Psi_2)$.

Proposition 2.5. *Let Φ be a C^2 topologically transitive Anosov flow on a closed Riemannian manifold M , $TM = T\Phi \oplus E^{ss} \oplus E^{uu}$ the Anosov splitting of Φ , and α^u a cocycle defined by $\alpha^u(p, t) = \log \det(D\Phi^t|_{E^{uu}(p)})$. Then, $P(\Phi, -\alpha^u) = 0$.*

Proof. See e.g. [8]. □

Corollary 2.6. *Let Φ be a C^2 Anosov flow with the Anosov splitting $TM = T\Phi \oplus E^{ss} \oplus E^{uu}$. Suppose that there exists $\lambda > 0$ such that*

$$J^u(p; \Phi) = \lambda \cdot \tau(p; \Phi) \tag{6}$$

for any $p \in \text{Per}(\Phi)$. Then, $h_{top}(\Phi) = \lambda$.

Proof. It is known that E^{uu} is Hölder continuous. Let α^u be the cocycle in the above proposition. By the Livschitz theorem, there exists a Hölder continuous function β such that $\alpha^u(p, t) = \lambda \cdot t + \beta(p) - \beta \circ \Phi^t(p)$. By Lemma 2.4 and Proposition 2.5, we have

$$h_{top}(\Phi) = P(0, \Phi) = P(-\alpha^u, \Phi) + \lambda = \lambda.$$

□

¹ Usually, we define the topological pressure for a function using its integral along the orbits of the flow (see e.g. [8]). Hence, if a cocycle α is differentiable with respect to t , then our definition is equivalent to the usual one for the function $(\partial\alpha/\partial t)(\cdot, 0)$.

2.3 Cocompact lattices of $\widetilde{\mathrm{PSL}}(2, \mathbb{R})$

Let $\mathrm{PSL}(2, \mathbb{R})$ be the quotient group of the special linear group of $\mathbb{R}^2$ by its center $\{\pm I\}$ and $\widetilde{\mathrm{PSL}}(2, \mathbb{R})$ the universal covering group of $\mathrm{PSL}(2, \mathbb{R})$. By θ , we denote the projection from $\widetilde{\mathrm{PSL}}(2, \mathbb{R})$ to $\mathrm{PSL}(2, \mathbb{R})$. It is known that the center of $\widetilde{\mathrm{PSL}}(2, \mathbb{R})$ coincides with the kernel of the projection θ and it is a infinite cyclic group. An element of $\widetilde{\mathrm{PSL}}(2, \mathbb{R})$ is called *central* if it is contained in the center of $\widetilde{\mathrm{PSL}}(2, \mathbb{R})$.

We define one-parameter subgroups $\{X^t\}_{t \in \mathbb{R}}$, $\{S^x\}_{x \in \mathbb{R}}$, and $\{U^y\}_{y \in \mathbb{R}}$ of $\widetilde{\mathrm{PSL}}(2, \mathbb{R})$ by

$$\theta(X^t) = \begin{bmatrix} e^{\frac{t}{2}} & 0 \\ 0 & e^{-\frac{t}{2}} \end{bmatrix}, \quad \theta(S^x) = \begin{bmatrix} 1 & x \\ 0 & 1 \end{bmatrix}, \quad \theta(U^y) = \begin{bmatrix} 1 & 0 \\ y & 1 \end{bmatrix}. \quad (7)$$

They are well-defined and satisfy the following identities:

$$S^x X^t = X^t S^{e^{-t}x}, \quad (8)$$

$$U^y X^t = X^t U^{e^t y}. \quad (9)$$

We define the trace $\mathrm{tr} P$ of each $P \in \widetilde{\mathrm{PSL}}(2, \mathbb{R})$ by the lift of the usual trace for matrices to $\widetilde{\mathrm{PSL}}(2, \mathbb{R})$. A non-central element P of $\widetilde{\mathrm{PSL}}(2, \mathbb{R})$ is called *hyperbolic*, *elliptic*, or *parabolic*, if $|\mathrm{tr} P|$ is greater, less than, or equal to two, respectively. If P is hyperbolic, then $\theta(P)$ is diagonalizable. Hence, there exists $Q \in \widetilde{\mathrm{PSL}}(2, \mathbb{R})$ and a central element $c \in \mathrm{PSL}(2, \mathbb{R})$ such that $P = cQX^{L(P)}Q^{-1}$. It implies that

$$|\mathrm{tr} P| = |\mathrm{tr} X^{L(P)}| = e^{\frac{L(P)}{2}} + e^{-\frac{L(P)}{2}}. \quad (10)$$

We summarize basic properties of cocompact lattices of $\widetilde{\mathrm{PSL}}(2, \mathbb{R})$. By $[P, Q]$ we denote the commutator $P^{-1}Q^{-1}PQ$ of $P, Q \in \widetilde{\mathrm{PSL}}(2, \mathbb{R})$.

Proposition 2.7. *Any cocompact lattice Γ of $\widetilde{\mathrm{PSL}}(2, \mathbb{R})$ satisfies the following properties:*

1. *If $P \in \widetilde{\mathrm{PSL}}(2, \mathbb{R})$ commutes with any element of Γ , then P is central.*
2. *$\theta(Q)$ is of finite order for any elliptic element $Q \in \widetilde{\mathrm{PSL}}(2, \mathbb{R})$.*
3. *Γ contains no parabolic element.*
4. *The commutator subgroup $[\Gamma, \Gamma]$ contains a non-trivial central element.*

Proof. We start with the proof of the first assertion. Consider the natural left action of $\mathrm{PSL}(2, \mathbb{R})$ on the real projective line $\mathbb{RP}^1$. Put $F(\gamma) = \{p \in \mathbb{RP}^1 \mid \gamma \cdot p = p\}$. Since Γ is cocompact, it contains two hyperbolic elements γ_1 and γ_2 such that $F(\gamma_1) \neq F(\gamma_2)$. It is easy to see that any element of $\widetilde{\mathrm{PSL}}(2, \mathbb{R})$ commuting with γ_i fixes each point of $F(\gamma_i)$. Hence, if γ_c commutes with all

element of Γ , then $F(\gamma_c)$ contains at least four points. It implies that γ_c is central since any non-central element fixes at most two points.

Next, we show that $\theta(\Gamma)$ is a cocompact lattice of $\widetilde{\mathrm{PSL}}(2, \mathbb{R})$. Let e be the identity element of $\widetilde{\mathrm{PSL}}(2, \mathbb{R})$ and G_c be the center of $\widetilde{\mathrm{PSL}}(2, \mathbb{R})$. Since Γ is cocompact, there exists a generator $\{\gamma_1, \dots, \gamma_m\}$ of Γ . Take two neighborhoods U and V of e such that $U \cap G_c = U \cap \Gamma = \{e\}$, $V \subset U$, and $[\gamma, \gamma_i] \in U$ for any $\gamma \in V$ and $i = 1, \dots, m$. If $\gamma \in \Gamma \cap cV$ for some central element c of $\widetilde{\mathrm{PSL}}(2, \mathbb{R})$, then

$$[\gamma, \gamma_i] = [c^{-1}\gamma, \gamma_i] \in U \cap \Gamma = \{e\}.$$

By the first assertion, we have $\gamma \in G_c$, and hence, $\gamma = c$. It implies that $\theta(\Gamma)$ is discrete. Since Γ is cocompact, also $\theta(\Gamma)$ is.

If γ is an elliptic element of $\widetilde{\mathrm{PSL}}(2, \mathbb{R})$, the closure of $\{\theta(\gamma)^n \mid n \in \mathbb{Z}\}$ is compact. It implies the second assertion. It is known that any cocompact lattice of $\mathrm{PSL}(2, \mathbb{R})$ contains no parabolic elements (see *e.g.* [2]). The third assertion follows from this fact.

To show the last assertion, we recall Selberg's lemma for cocompact lattices of $\mathrm{PSL}(2, \mathbb{R})$. It asserts the existence of a finite index subgroup Γ' of $\theta(\Gamma)$ which contains no elliptic element. Since Γ' is a cocompact lattice and any non-identity element is hyperbolic, the quotient map $\pi : \mathrm{PSL}(2, \mathbb{R}) \rightarrow \Gamma' \backslash \mathrm{PSL}(2, \mathbb{R})$ gives the unit tangent bundle over a closed surface with negative Euler number. It implies that the commutator subgroup of $\theta^{-1}(\Gamma') \cap \Gamma$ contains a non-trivial central element. $\square$

The following lemma gives an estimate of the translation length of a commutator.

Lemma 2.8. *Let P and Q be elements of $\widetilde{\mathrm{PSL}}(2, \mathbb{R})$ such that P is hyperbolic and $[P, Q]$ is neither central nor parabolic. Then,*

$$\lim_{n \rightarrow \infty} \frac{L([P^n, Q])}{2n} = L(P)$$

Proof. Take $A \in \widetilde{\mathrm{PSL}}(2, \mathbb{R})$ and a central element C so that $P = A^{-1}X^{L(P)}AC$. Put

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} = \theta(AQA^{-1}).$$

Then, we have

$$\theta(A[P^n, Q]A^{-1}) = \begin{bmatrix} ad - e^{-nL(P)}bc & (1 - e^{-nL(P)})bd \\ (1 - e^{nL(P)})ac & ad - e^{nL(P)}bc \end{bmatrix}.$$

The equation for $n = 1$ implies that $bc \neq 0$ since $[P, Q]$ is neither central or

parabolic. Now, we have

$$\begin{aligned}
\lim_{n \rightarrow \infty} \frac{L([P^n, Q])}{2n} - L(P) &= \lim_{n \rightarrow \infty} \frac{1}{n} (\log |\operatorname{tr} [P^n, Q]| - \log |\operatorname{tr} P^{2n}|) \\
&= \lim_{n \rightarrow \infty} \frac{1}{n} (\log |\operatorname{tr} (A[P^n, Q]A^{-1})| - \log |\operatorname{tr} P^{2n}|) \\
&= \lim_{n \rightarrow \infty} \frac{1}{n} \log \left| \frac{2ad - (bc)\operatorname{tr} P^{2n}}{\operatorname{tr} P^{2n}} \right| \\
&= \lim_{n \rightarrow \infty} \frac{bc}{n} = 0.
\end{aligned}$$

□

Let Γ be a cocompact lattice of $\widetilde{\operatorname{PSL}}(2, \mathbb{R})$. By M_Γ , we denote the quotient space $\Gamma \backslash \widetilde{\operatorname{PSL}}(2, \mathbb{R})$. We identify the lattice Γ with the fundamental group of M_Γ . The following is a direct consequence of Proposition 2.7

Lemma 2.9. *Any non-hyperbolic element of Γ represents the zero class of $H_1(M_\Gamma, \mathbb{R})$.* □

Lemma 2.10. *For any $\gamma \in \Gamma$ with $L(\gamma) > 0$, there exist $P \in \widetilde{\operatorname{PSL}}(2, \mathbb{R})$, a central element $c \in \Gamma$, and an integer $n \geq 1$ such that $(c\gamma^n)P = PX^{nL(\gamma)}$.*

Proof. Since any hyperbolic element of $\operatorname{PSL}(2, \mathbb{R})$ is diagonalizable, there exists $P \in \operatorname{PSL}(2, \mathbb{R})$ and a central element $c_0 \in \widetilde{\operatorname{PSL}}(2, \mathbb{R})$ such that $c_0\gamma P = PX^{L(\gamma)}$. Since the center of $\widetilde{\operatorname{PSL}}(2, \mathbb{R})$ is cyclic, the last assertion of Proposition 2.7 implies that $c = c_0^n$ is contained in Γ for some $n \geq 1$. Now, it is easy to check that $(c\gamma^n)P = PX^{nL(\gamma)}$. □

3 Anosov flows associated to actions

In this section, we study the relation between the period of a periodic points, expansion along invariant subbundles, and the one-form ω in the equation (2). It leads to the proof of an easy half of Theorem B (Subsection 3.4) and Theorem A (Subsection 3.5).

3.1 Homogeneous Anosov flows

Fix a cocompact lattice Γ of $\widetilde{\operatorname{PSL}}(2, \mathbb{R})$. Let X_Γ , Y_Γ^s , and Y_Γ^u , be the vector fields on M_Γ which generate the natural right action of the one-parameter subgroups $\{X^t\}_{t \in \mathbb{R}}$, $\{S^x\}_{x \in \mathbb{R}}$, and $\{U^y\}_{y \in \mathbb{R}}$ of $\operatorname{PSL}(2, \mathbb{R})$ on $M_\Gamma = \Gamma \backslash \widetilde{\operatorname{PSL}}(2, \mathbb{R})$ from right. Then, they satisfy the identities

$$[Y_\Gamma^u, X_\Gamma] = Y_\Gamma^u, \quad [Y_\Gamma^s, X_\Gamma] = -Y_\Gamma^s. \quad (11)$$

In the below, we identify GA with a subgroup $\{X^t U^y \mid t, y \in \mathbb{R}\}$ of $\widetilde{\operatorname{PSL}}(2, \mathbb{R})$. We define the homogeneous action $\rho_\Gamma : M_\Gamma \times \text{GA} \rightarrow M_\Gamma$ associated to Γ by

$$\rho_\Gamma(\Gamma P, X^t U^y) = \Gamma(PX^t U^y)$$

for $P \in \widetilde{\text{PSL}}(2, \mathbb{R})$ and $X^t U^y \in \text{GA}$. We also define the flow Φ_Γ on M_Γ by $\Phi_\Gamma^t(\Gamma P) = \Gamma(PX^t)$. It is generated by X_Γ . For each periodic point p of Φ_Γ , let $\gamma_\Gamma(p)$ be the element of Γ such that

$$\gamma_\Gamma(p)P = PX^{\tau(p, \Phi_\Gamma)}, \quad (12)$$

where $P \in \theta^{-1}(p)$.

The uniqueness of the Anosov splitting and the equations (11) and (12) imply that Φ_Γ is an Anosov flow with the Anosov splitting $TM = T\Phi_\Gamma \oplus \mathbb{R} \cdot Y_\Gamma^u \oplus \mathbb{R} \cdot Y_\Gamma^s$, and it satisfies

$$J^u(p; \Phi_\Gamma) = -J^s(p; \Phi_\Gamma) = \tau(p; \Phi_\Gamma) = L(\gamma_\Gamma(p)). \quad (13)$$

By $\mathcal{F}_\Gamma^u$, we denote the weak unstable foliation of Φ_Γ . It coincides with the orbit foliation of the natural right action ρ_Γ of GA on M_Γ .

Lemma 3.1. *$H_1(M_\Gamma, \mathbb{R})$ is generated by periodic orbits of Φ_Γ as an $\mathbb{R}$ -vector space.*

Proof. Lemma 2.9 implies that any non-zero integer class c is represented by a hyperbolic element of Γ . By Lemma 2.10, there exists $n \geq 1$ such that $n \cdot c$ is represented by a periodic orbit of Φ_Γ . $\square$

Lemma 3.2. *There exists a periodic orbit of Φ_Γ which represents the zero class of $H^1(M_\Gamma)$.*

Proof. It is easy to see that $[\Gamma, \Gamma]$ contains a hyperbolic element. By Lemma 2.10, the zero class is represented by a periodic orbit of Φ_Γ . $\square$

Lemma 3.3. *The flow Φ_Γ is topologically transitive.*

Proof. It is known that any volume-preserving Anosov flow is topologically transitive. Since Φ_Γ preserves the smooth volume induced from the Haar measure on $\widetilde{\text{PSL}}(2, \mathbb{R})$, it is topologically transitive. $\square$

The following proposition characterizes Anosov flows whose weak unstable foliation is $\mathcal{F}_\Gamma^u$.

Proposition 3.4. *Let Φ be an Anosov flow on M_Γ such that its weak unstable foliation coincides with $\mathcal{F}_\Gamma^u$. Then, there exists a topological equivalence between Φ_Γ and Φ which is isotopic to the identity and preserves each leaf of $\mathcal{F}_\Gamma^u$.*

Proof. In [10], Ghys showed that any Anosov flows on the unit tangent bundle of a closed surface with a hyperbolic metric is topologically equivalent to the geodesic flow. A careful application of his argument gives a proof of the proposition. The key observation for our case is the following: The assumption on the weak unstable foliation implies that the asymptotics of the lifts of the periodic orbits of Φ and Φ_Γ to $\widetilde{\text{PSL}}(2, \mathbb{R})$ coincide if and only if they are contained in the same leaf of the lift of $\mathcal{F}_\Gamma^u$. $\square$

Let ρ be an action in $\mathcal{A}_\Gamma$. We define a flow Φ_ρ on M_Γ by $\Phi_\rho^t(p) = \rho(p, X^t)$. By Theorem 1.5, it is an Anosov flow. The weak unstable foliation of Φ_ρ coincides with the orbit foliation of ρ , and hence, with $\mathcal{F}_\Gamma^u$. Let H_ρ be the topological equivalence which is obtained by the above proposition. By Lemmas 3.1 and 3.2, Φ_ρ is topologically transitive and it admits a periodic point p whose orbit is null-homologous.

Let Y_ρ^u be the vector field which generates the ρ -action of $\{U^y\}_{y \in \mathbb{R}}$. Since $D\Phi_\rho^t(Y_\rho^u) = e^t \cdot Y_\rho^u$, the vector field Y_ρ^u is tangent to the strong unstable foliation and

$$J^u(q; \Phi_\rho) = \tau(q; \Phi_\rho) \quad (14)$$

for any $q \in \text{Per}(\Phi_\rho) = H_\rho(\text{Per}(\Phi_\Gamma))$. For each $p \in \text{Per}(\Phi_\Gamma)$, the Φ_Γ -orbit of p is homotopic to the Φ_ρ -orbit of $H_\rho(p)$ in $\mathcal{F}_\Gamma^u(p)$. It implies that

$$J^s(H_\rho(p); \Phi_\rho) = J^s(p; \Phi_\Gamma) = -L(\gamma_\Gamma(p)) = -\tau(p; \Phi_\Gamma). \quad (15)$$

In particular,

$$J(H_\rho(p); \Phi_\rho) = \tau(H_\rho(p); \Phi_\rho) - \tau(p; \Phi_\Gamma). \quad (16)$$

3.2 Reparametrization of Φ_Γ

For $a \in H^1(M_\Gamma, \mathbb{R})$, we define a left-action Γ_a of Γ on $\widetilde{\text{PSL}}(2, \mathbb{R})$ by

$$\Gamma_a(\gamma, P) = \gamma P X^{a(\gamma)}.$$

This action was introduced by Ghys in [12], where he used it to construct a non-homogeneous Anosov flow with a smooth Anosov splitting.

The following proposition is due to Salein [25]. However, we give a proof here since the presentation of the statement is different from Salein's and the proof is short.

Proposition 3.5. *For any $a \in \Delta_\Gamma$, the action Γ_a is free and properly discontinuous.*

Proof. Since $a \in \Delta_\Gamma$, there exists $\epsilon > 0$ such that $|a(\gamma)| < (1 - \epsilon)L(\gamma)$ for any $\gamma \in \Gamma$ with $L(\gamma) \neq 0$.

First, we show that the action is free. If $\Gamma_a(\gamma, P) = P$, then $\gamma = P X^{-a(\gamma)} P^{-1}$. It implies $L(\gamma) = |a(\gamma)|$. By the assumption, we have $a(\gamma) = 0$, and hence, γ is the identity.

To show that the action is properly discontinuous, we fix a compact subset V of $\widetilde{\text{PSL}}(2, \mathbb{R})$. For $K > 0$, put $\Gamma(K) = \{\gamma \in \Gamma \mid L(\gamma) \leq K\}$. We claim that the set $\{\gamma \in \Gamma(K) \mid \Gamma_a(\gamma \times V) \cap V \neq \emptyset\}$ is finite for any $K > 0$. Let d be a left invariant distance on $\widetilde{\text{PSL}}(2, \mathbb{R})$ and e the identity element of $\widetilde{\text{PSL}}(2, \mathbb{R})$. Put $\delta_K = \sup\{d(X^t, e) \mid |t| \leq K\}$ and $V_K = \{P \in \widetilde{\text{PSL}}(2, \mathbb{R}) \mid \inf_{Q \in V} d(P, Q) \leq \delta_K\}$. Since

$$d(\Gamma_a(\gamma, P), \gamma \cdot P) = d(X^{a(\gamma)}, e) \leq \delta_K$$

for any $P \in V$ and $\gamma \in \Gamma(K)$, we have $\Gamma_a(\gamma \times V) \subset \gamma \cdot V_K$ for any $\gamma \in \Gamma(K)$. Since the natural left action of Γ on $\widetilde{\text{PSL}}(2, \mathbb{R})$ is properly discontinuous, the set $\{\gamma \in \Gamma(K) \mid \gamma \cdot V_K \cap V_K \neq \emptyset\}$ is finite. It implies the claim.

Now, it is sufficient to show that there exists $K > 0$ such that $\Gamma_a(\gamma \times V) \cap V = \emptyset$ for any $\gamma \in \Gamma(K)$. Since $\widetilde{\text{PSL}}(2, \mathbb{R})$ is the universal covering group of the special linear group of $\mathbb{R}^2$, it acts on $\mathbb{R}^2$ naturally as linear transformations. Let $\|A\|$ be the operator norm of $A \in \text{PSL}(2, \mathbb{R})$ with respect to the Euclid norm on $\mathbb{R}^2$. It is easy to see that $L(P) \leq 2 \log \|P\|$ for any $P \in \widetilde{\text{PSL}}(2, \mathbb{R})$ and $|t| = L(X^t) = 2 \log \|X^t\|$ for any $t \in \mathbb{R}$. Put $K_0 = \sup_{P, Q \in V} \log \|P^{-1}Q\|$. Then, any $P, Q \in V$ and $\gamma \in \Gamma(2K_0\epsilon^{-1})$ satisfy

$$\begin{aligned} |P^{-1}QX^{-a(\gamma)}| &\leq 2 \log(\|P^{-1}Q\| \cdot \|X^{-a(\gamma)}\|) \\ &\leq 2K_0 + |a(\gamma)| < \epsilon L(\gamma) + |a(\gamma)| < L(\gamma), \end{aligned}$$

Since $L(P^{-1} \cdot \Gamma_a(\gamma, P) \cdot X^{-a(\gamma)}) = L(P^{-1}\gamma P) = L(\gamma)$, we have $\Gamma_a(\gamma, P) \neq Q$. It implies that $\Gamma_a(\gamma) \cdot V \cap V = \emptyset$ for any $\gamma \in \Gamma(2K_0\epsilon^{-1})$. $\square$

Take $a \in \Delta_\Gamma$. Put $M_{\Gamma, a}$ be the quotient space of $\widetilde{\text{PSL}}(2\mathbb{R})$ by the action of Γ_a . Let $\tilde{\Phi}_\Gamma$ be the flow on $\widetilde{\text{PSL}}(2, \mathbb{R})$ defined by $\tilde{\Phi}_\Gamma(P) = P \cdot X^t$, and $\tilde{\mathcal{F}}^{ss}$ and $\tilde{\mathcal{F}}^{uu}$ be C^ω foliations defined by $\tilde{\mathcal{F}}^{ss}(P) = \{PS^x \mid x \in \mathbb{R}\}$ and $\tilde{\mathcal{F}}^{uu}(P) = \{PU^y \mid y \in \mathbb{R}\}$. The flow $\tilde{\Phi}_\Gamma$ commutes with the action of Γ_a and preserves foliations $\tilde{\mathcal{F}}^{ss}$ and $\tilde{\mathcal{F}}^{uu}$. Hence, $\tilde{\Phi}_\Gamma$ induces a C^ω flow $\Phi_{\Gamma, a}$ on $M_{\Gamma, a}$ and the foliations $\tilde{\mathcal{F}}^{ss}$ and $\tilde{\mathcal{F}}^{uu}$ project to C^ω $\Phi_{\Gamma, a}$ -invariant foliations $\mathcal{F}_a^{ss}$ and $\mathcal{F}_a^{uu}$. It is easy to see that $\Phi_{\Gamma, a}$ is an Anosov flow with the C^ω Anosov splitting $TM_{\Gamma, a} = T\Phi_{\Gamma, a} \oplus T\mathcal{F}_a^{ss} \oplus T\mathcal{F}_a^{uu}$.

Ghys showed that the Φ_a can be identified with a time-change of Φ_Γ .

Proposition 3.6 ([11]). *There exists a C^ω equivalence H_a between Φ_Γ and $\Phi_{\Gamma, a}$ which is isotopic to the identity and satisfies*

$$\tau(H_a(p); \Phi_{\Gamma, a}) = \tau(p; \Phi_\Gamma) + a(\gamma_\Gamma(p)) \quad (17)$$

for any $p \in \text{Per}(\Phi_\Gamma)$.

3.3 The cohomology class $\bar{a}(\rho)$

Take $\rho \in \mathcal{A}_\Gamma$ and a closed one-form ω satisfying (2). Let X_ρ be the vector field generating Φ_ρ .

Proposition 3.7. *The one-form ω is uniquely determined.*

Proof. Suppose that closed one forms ω and ω' satisfy the equation (2). Then, $(\Phi_\rho^t)^*(\omega - \omega') = \omega - \omega'$ for any $t \in \mathbb{R}$. In other words, $\omega - \omega'$ is Φ_ρ -invariant. Since $\lim_{t \rightarrow \infty} \|D\Phi_\rho^t|_{E_\rho^{ss}}\| = 0$, we have

$$(\omega - \omega')(v^s) = \lim_{t \rightarrow \infty} (\Phi_\rho^t)^*(\omega - \omega')(v^s) = \lim_{t \rightarrow \infty} (\omega - \omega')(D\Phi_\rho^t(v^s)) = 0$$

for any $v^s \in E_\rho^{ss}$. A similar argument shows that $(\omega - \omega')(v^u) = 0$ for any $v^u \in E_\rho^{uu}$.

The Φ_ρ -invariance of $(\omega - \omega')$ implies $(\omega - \omega')(X_\rho)$ is a Φ_ρ -invariant function. Since Φ_ρ is topologically transitive, $(\omega - \omega')(X_\rho)$ is a constant function. Take a null-homologous periodic orbit γ of Φ_ρ . Then, the integral of $\omega - \omega'$ along γ is zero. It implies that $(\omega - \omega')(X_\rho) = 0$. $\square$

Recall that $\bar{a}(\rho)$ is the cohomology class of ω . We define a function $\bar{\delta}$ on $\mathcal{A}_\Gamma$ by

$$\bar{\delta}(\rho) = 1 - (h_{top}(\Phi_{\Gamma, \bar{a}(\rho)}))^{-1}.$$

It depends only on $\bar{a}(\rho)$.

Proposition 3.8.

$$\log \det D\Phi_\rho^t(p) = \bar{\delta}(\rho) \cdot t + \int_{\{\Phi_\rho^s(p)\}_{0 \leq s \leq t}} \omega \quad (18)$$

for any $(p, t) \in M_\Gamma \times \mathbb{R}$.

Proof. Put $f(p) = \frac{d}{dt}(\log \det D\Phi_\rho^t(p))|_{t=0}$. By the equation (2), we have

$$\begin{aligned} \omega(X_\rho) \circ \Phi^t - \omega(X_\rho) &= (\Phi_\rho^* \omega)(X_\rho) - \omega(X_\rho) \\ &= d(\log \det D\Phi_\rho^t)(X_\rho) \\ &= f \circ \Phi^t - f. \end{aligned}$$

In particular, the function $\omega(X_\rho) - f$ is Φ_ρ -invariant. Since Φ_ρ is topologically transitive, $\omega(X_\rho) - f$ is a constant function. Hence, there exists a real number δ satisfying the equation (18).

Take a periodic point q of Φ_ρ whose orbit is null-homologous. Then, we have

$$\delta \cdot \tau(q; \Phi_\rho) = J(q; \Phi_\rho) < J^u(q; \Phi_\rho) = \tau(q; \Phi_\rho)$$

It implies that $\delta < 1$. By the equations (14), (15), and (18), we have

$$(1 - \delta)\tau(H_\rho(p); \Phi_\rho) = \tau(p; \Phi_\Gamma) + \bar{a}(\rho)(\gamma_\Gamma(p)) \quad (19)$$

for any $p \in \text{Per}(\Phi_\Gamma)$.

We show that $\delta = \bar{\delta}(\rho)$. Let Ψ be an Anosov flow on M_Γ defined by $\Psi^t = \Phi_\rho^{\frac{t}{1-\delta}}$. Then, $\text{Per}(\Psi) = \text{Per}(\Phi_\rho)$ and any $p \in \text{Per}(\Psi)$ satisfies $J^u(p; \Psi) = J^u(p; \Phi_\rho)$ and

$$\tau(p; \Psi) = (1 - \delta)\tau(p; \Phi_\rho). \quad (20)$$

In particular, $J^u(p; \Psi) = (1 - \delta)^{-1}\tau(p; \Psi)$. By Corollary 2.6, we have $h_{top}(\Psi) = (1 - \delta)^{-1}$.

By (17), (19), and (20), $H_\rho \circ H_{\bar{a}(\rho)}^{-1}$ is a topological equivalence between $\Phi_{\Gamma, \bar{a}(\rho)}$ and Ψ which preserves the period of each periodic point. Hence, Theorem 2.2 implies that Ψ and $\Phi_{\Gamma, \bar{a}(\rho)}$ are topologically conjugate. Therefore, we have $h_{top}(\Phi_{\Gamma, \bar{a}(\rho)}) = h_{top}(\Psi) = (1 - \delta)^{-1}$. $\square$

3.4 The space Δ_Γ

First, we rewrite Δ_Γ in terms of Φ_Γ .

Lemma 3.9.

$$\Delta_\Gamma = \left\{ a \in H^1(M_\Gamma, \mathbb{R}) \mid \inf_{p \in \text{Per}(\Phi_\Gamma)} \frac{a(\gamma_\Gamma(p))}{\tau(p; \Phi_\Gamma)} > -1 \right\}. \quad (21)$$

Proof. Fix $a \in H^1(M_\Gamma, \mathbb{R})$ and $\gamma \in \Gamma$ with $L(\gamma) \neq 0$. Applying Lemma 2.10 to γ and γ^{-1} , we obtain $p, p' \in \text{Per}(\Phi_\Gamma)$, $n, n' \geq 1$, and central elements $c, c' \in \Gamma$ such that $\gamma_\Gamma(p) = c\gamma^n$ and $\gamma_\Gamma(p') = c'\gamma^{-n'}$. Lemma 2.9 and the equation (13), we have

$$\frac{a(\gamma)}{L(\gamma)} = \frac{a(\gamma_\Gamma(p))}{\tau(p; \Phi_\Gamma)} = -\frac{a(\gamma_\Gamma(p'))}{\tau(p'; \Phi_\Gamma)}.$$

□

The following is an easy half of Theorem B.

Proposition 3.10. $\bar{a}(\rho) \in \Delta_\Gamma$ for any $\rho \in \mathcal{A}_\Gamma$.

Proof. Take $\rho \in \mathcal{A}_\Gamma$. Since Φ_ρ is an Anosov flow on a closed manifold M_Γ , there exists $C > 0$ such that

$$-\log \|D\Phi_\rho^t|_{E^{ss}(q)}\| \leq C(1 - \bar{\delta}(\rho))t \quad (22)$$

for any $q \in M_\Gamma$ and $t \geq 0$. It implies $|J^s(q, \Phi_\rho)| \leq C(1 - \bar{\delta}(\rho))\tau(q; \Phi_\rho)$ for any $q \in \text{Per}(\Phi_\rho)$. By (15) and (19), we have

$$1 + \frac{\bar{a}(\rho)(\gamma_\Gamma(p))}{\tau(p; \Phi_\Gamma)} = \frac{(1 - \bar{\delta}(\rho)) \cdot \tau(H_\rho(p); \Phi_\rho)}{|J^s(H_\rho(p); \Phi_\rho)|} \geq \frac{1}{C}$$

for any $p \in \text{Per}(\Phi_\Gamma)$. The equation (21) implies that $\bar{a}(\rho)$ is an element of Δ_Γ . □

Recall that X_Γ is the vector field generating Φ_Γ . The following proposition gives another characterization of Δ_Γ .

Proposition 3.11. A cohomology class $a \in H^1(M_\Gamma, \mathbb{R})$ is contained in Δ_Γ if and only if it is represented by a closed one-form ω on M_Γ satisfying

$$\inf_{p \in M_\Gamma} \omega(X_\Gamma)(p) > -1.$$

Corollary 3.12. Δ_Γ is a non-empty, open, and convex subset of $H^1(M_\Gamma, \mathbb{R})$. □

Proof of Proposition 3.11. First, we suppose that a is represented by a closed one form ω such that $\inf_{p \in M_\Gamma} \omega(X_\Gamma)(p) = -1 + \epsilon$ for some $\epsilon > 0$. Then, we have $a(\gamma_\Gamma(p)) \geq (-1 + \epsilon)\tau(p; \Phi_\Gamma)$ for any $p \in \text{Per}(\Phi_\Gamma)$. By Lemma 3.9, a is an element of Δ_Γ .

Next, we suppose that a is contained in Δ_Γ . Then, there exists $\epsilon' > 0$ such that $a(\gamma_\Gamma(p)) \geq (-1 + \epsilon')\tau(p; \Phi_\Gamma)$ for any $p \in \text{Per}(\Phi_\Gamma)$. By Proposition 2.3, there exists a C^∞ function β such that $\inf_{p \in M_\Gamma} (\omega + du)(X_\Gamma)(p) \geq -1 + \epsilon'/2$. □

3.5 Proof of Theorem A

Now, we prove Theorem A. Let Γ be a cocompact lattice of $\widetilde{\text{PSL}}(2, \mathbb{R})$. Suppose that two actions $\rho_1, \rho_2 \in \mathcal{A}_\Gamma$ satisfy $\bar{a}_\Gamma(\rho_1) = \bar{a}_\Gamma(\rho_2) = a$. Put $\delta = 1 - h_{top}(\Phi_{\Gamma, a})^{-1}$. Then, $\bar{\delta}(\rho_1) = \bar{\delta}(\rho_2) = \delta$ by definition. The equation (19) implies

$$\tau(H_{\rho_1}(p); \Phi_{\rho_1}) = \tau(H_{\rho_2}(p); \Phi_{\rho_2}) = (1 - \delta)^{-1} \{ \tau(p; \Phi_\Gamma) + a(\gamma_\Gamma(p)) \}$$

for any $p \in \text{Per}(\Phi_\Gamma)$. By Theorem 2.2, there exists a continuous function β such that the map H defined by $H(p) = \Phi_{\rho_2}^{\beta(p)} \circ H_{\rho_2} \circ H_{\rho_1}^{-1}$ is a topological conjugacy between Φ_{ρ_1} and Φ_{ρ_2} . By (14) and (15),

$$\begin{aligned} J^s(H_{\rho_1}(p); \Phi_{\rho_1}) &= J^s(H_{\rho_2}(p); \Phi_{\rho_2}) = -\tau(p; \Phi_\Gamma), \\ J^u(H_{\rho_1}(p); \Phi_{\rho_1}) &= J^u(H_{\rho_2}(p); \Phi_{\rho_2}) = (1 - \delta)^{-1} \{ \tau(p; \Phi_\Gamma) + a(\gamma_\Gamma(p)) \} \end{aligned}$$

for any $p \in \text{Per}(\Phi_\Gamma)$. By Theorem 2.1, the conjugacy H is a C^∞ diffeomorphism.

Let X_{ρ_i} be the vector field generating Φ_{ρ_i} and U_{ρ_i} be the vector field generating the flow $\{\rho_i(\cdot, U^y)\}_{y \in \mathbb{R}}$ for each $i = 1, 2$. Since $[U_{\rho_i}, X_{\rho_i}] = U_{\rho_i}$, the vector field U_{ρ_i} is tangent to the strong unstable foliation of Φ_{ρ_i} . Since the C^∞ conjugacy H preserves the strong unstable foliation, there exists a C^∞ function g on M_Γ such that $g(p) \neq 0$ and

$$DH(U_{\rho_1})(p) = g(p) \cdot U_{\rho_2}(p)$$

for any $p \in M_\Gamma$. The equations $[U_{\rho_i}, X_{\rho_i}] = U_{\rho_i}$ and $DH(X_{\rho_1}) = X_{\rho_2}$ imply $X_{\rho_2}g = 0$. Since the flow Φ_{ρ_2} is topologically transitive, the function g is constant with a non-zero value λ . Now, we have

$$\rho_2(H(p), X^t U^{\lambda y}) = H(\rho_1(p, X^t U^y))$$

for any $t, y \in \mathbb{R}$ and $p \in M_\Gamma$. It implies that ρ_1 and ρ_2 are C^∞ conjugate.

4 Construction of non-homogeneous actions

In this section, we prove a harder half of Theorem B and Theorem C, *i.e.*, construction of action $\rho \in \mathcal{A}_\Gamma$ which is C^∞ conjugate to an C^ω action and satisfies $\bar{a}_\Gamma(\rho) = a$ for any given $a \in \Delta_\Gamma$. It can be reduced to a construction of an C^ω Anosov flow with a special property (Proposition 4.3). The construction follows Cawley's method ([7]) on deformation of smooth structure of Anosov systems using measures along invariant foliations. In our case, we deform the Anosov flow $\Phi_{\Gamma, a}$, which is defined in Subsection 3.2, using the Margulis measure along the strong unstable foliation.

4.1 The Margulis measure

Let Φ be an Anosov flow on a closed manifold M and $TM = T\Phi \oplus E^{ss} \oplus E^{uu}$ its Anosov splitting. By $\mathcal{F}^{ss}$, $\mathcal{F}^{uu}$, $\mathcal{F}^s$, and $\mathcal{F}^u$, we denote the invariant foliations of Φ which is defined in Subsection 2.1.

For a foliation $\mathcal{F}$, we denote the leaf of $\mathcal{F}$ which contains p by $\mathcal{F}(p)$. For any $p, q \in M$ which are sufficiently close to each other, we can define the holonomy map $\tilde{h}_{pq}$ of $\mathcal{F}^{ss}$ from a neighborhood $\tilde{U}_{pq}$ of p in $\mathcal{F}^u(p)$ to $\mathcal{F}^u(q)$.

Theorem 4.1 (Margulis [22]). *There exists a family $\{\tilde{\mu}_p\}_{p \in M}$ which has the following properties:*

1. $\tilde{\mu}$ is a locally finite Borel measure on $\mathcal{F}^u(p)$ which is non-atomic and positive on each non-empty open subset of $\mathcal{F}^u(p)$.
2. If $\mathcal{F}^u(p) = \mathcal{F}^u(q)$, then $\tilde{\mu}_p = \tilde{\mu}_q$.
3. The family $\{\tilde{\mu}_p\}_{p \in M}$ is holonomy invariant, i.e., if p and q are sufficiently close to each other, then $\tilde{\mu}_q(\tilde{h}_{pq}(A)) = \tilde{\mu}_p(A)$ for any Borel subset A of $\tilde{U}_{pq}$.
4. There exists $\lambda > 0$ such that

$$\tilde{\mu}_p(\Phi^t(A)) = e^{\lambda t} \cdot \tilde{\mu}_p(A)$$

for any $p \in M$, $t \in \mathbb{R}$, and any Borel subset A of $\mathcal{F}^u(p)$.

As explained in Chapter 3 of [22], the family $\{\tilde{\mu}_p\}_{p \in M}$ induces a family $\{\mu_p\}_{p \in M}$ of measures on leaves of $\mathcal{F}^{uu}$ such that

$$\tilde{\mu}_p(\{\Phi^t(q) \mid q \in V, 0 \leq t \leq \epsilon\}) = \mu_p(V) \int_0^\epsilon e^{\lambda t} dt.$$

for any small open set $V \subset \mathcal{F}^{uu}(p)$ and any small $\epsilon > 0$. We call the family $\{\mu_p\}_{p \in M}$ the Margulis measure of Φ along $\mathcal{F}^{uu}$. It satisfies the following properties:

- For any small open subset $\tilde{V} \subset \mathcal{F}^u(p)$ of the form

$$\tilde{V} = \{\Phi^t(p') \mid p' \in V, t \in (\eta_-(p'), \eta_+(p'))\},$$

where V is an open subset of $\mathcal{F}^{uu}(p)$, and η_- and η_+ are continuous functions on V ,

$$\tilde{\mu}_p(\tilde{V}) = \int_V \left(\int_{\eta_-(p')}^{\eta_+(p')} e^{\lambda t} dt \right) d\mu_p(p'). \quad (23)$$

- For any $p \in M$, $t \in \mathbb{R}$, and any small open subset V of $\mathcal{F}^{uu}(p)$,

$$\mu_{\Phi^t(p)}(\Phi^t(V)) = e^{\lambda t} \cdot \mu_p(V) \quad (24)$$

When two points p and q in M are sufficiently close to each other, we can define the holonomy map h_{pq} of $\mathcal{F}^s$ from a neighborhood U_{pq} of p in $\mathcal{F}^{uu}(p)$ to $\mathcal{F}^{uu}(q)$.

Proposition 4.2. *Suppose that the flow Φ and the foliations $\mathcal{F}^{ss}$, $\mathcal{F}^{uu}$, $\mathcal{F}^s$, $\mathcal{F}^u$ are real analytic. Then, for any given $p \in M$, the Radon-Nikodym derivative $\frac{d(\mu_q \circ h_{qp})}{d\mu_q}(q)$ depends real analytically on $q \in M$.*

Proof. Fix a neighborhood V^{uu} of p in $\mathcal{F}^{uu}(p)$. There exists a neighborhood U of p in M and a C^ω function η on U such that $\Phi^{\eta(q)} \circ \tilde{h}_{pq}(V^{uu})$ contains q . Remark that the function η satisfies

$$\Phi^{\eta(h_{pq}(p'))} \circ \tilde{h}_{pq}(p') = h_{pq}(p') \quad (25)$$

for any $p' \in \mathcal{F}^{uu}(p)$ sufficiently close to p .

Take a small neighborhood $V \subset V^{uu}$ of p in $\mathcal{F}^{uu}$ and a small positive number $\epsilon > 0$. Put $V_\epsilon = \{\Phi^t(p') \mid p' \in V, t \in (-\epsilon, \epsilon)\}$. Since Φ preserves the foliation $\mathcal{F}^{ss}$, we have $\tilde{h}_{pq} \circ \Phi^t = \Phi^t \circ \tilde{h}_{pq}$ for any $t \in (-\epsilon, \epsilon)$. It implies

$$\begin{aligned} \tilde{h}_{pq}(V_\epsilon) &= \{\Phi^t(q') \mid q' \in \tilde{h}_{pq}(V), t \in (-\epsilon, \epsilon)\} \\ &= \{\Phi^t(q') \mid q' \in h_{pq}(V), t \in (-\epsilon - \eta(q'), \epsilon - \eta(q'))\}, \end{aligned}$$

where the latter equation follows from (25). By the holonomy invariance of $\{\tilde{\mu}_p\}_{p \in M}$ and the equation (23), we have

$$\int_{h_{pq}(V)} \left(\int_{-\epsilon - \eta(q')}^{\epsilon - \eta(q')} e^{\lambda t} dt \right) d\mu_q(q') = \int_V \left(\int_{-\epsilon}^{\epsilon} e^{\lambda t} dt \right) d\mu_p. \quad (26)$$

It implies

$$\int_{h_{pq}(V)} \exp(-\lambda \cdot \eta(q')) d\mu_q(q') = \int_V d\mu_p.$$

Hence, we obtain

$$\frac{d(\mu_p \circ h_{qp})}{d\mu_q}(q) = \exp(-\lambda \cdot \eta(q)).$$

It is a C^ω function on U . □

4.2 Codimension-one Anosov flows with constant expansion

We construct a codimension-one Anosov flow with constant unstable expansion by a deformation preserving the strong unstable foliation.

Proposition 4.3. *Let Φ be a C^ω Anosov flow on a closed manifold M . Suppose that its Anosov splitting is of class C^ω and the strong unstable foliation $\mathcal{F}^{uu}$ is one-dimensional. Then, there exists another C^ω Anosov flow $\check{\Phi}$ on a C^ω closed manifold $\check{M}$, a C^ω vector field Y^u , a C^∞ diffeomorphism $\check{H} : M \rightarrow \check{M}$, and a constant $\lambda > 0$ such that Y^u generates the strong unstable foliation $\check{\mathcal{F}}^{uu}$ of $\check{\Phi}$ and*

$$\check{H} \circ \Phi^t(\mathcal{F}^{uu}(p)) = \check{\Phi}^t(\check{\mathcal{F}}^{uu}(\check{H}(p))), \quad (27)$$

$$D\check{\Phi}^t(Y^u(p)) = e^{\lambda t} \cdot Y^u(\check{\Phi}^t(p)) \quad (28)$$

for any $p \in M$ and $t \in \mathbb{R}$,

The equation (27) implies that $\check{H}$ maps the weak unstable foliation of Φ to that of $\check{\Phi}$. Since $\Phi^t(\mathcal{F}^{uu}(p)) = \mathcal{F}^{uu}(p)$ for $p \in \text{Per}(\Phi)$ if and only if t is an integer multiple of $\tau(p; \Phi)$, it also implies

$$\tau(\check{H}(p), \check{\Phi}) = \tau(p, \Phi) \quad (29)$$

for any $p \in \text{Per}(\Phi)$.

The rest of the subsection is devoted to the proof of the proposition. Let N be the dimension of the manifold M , $\mathcal{F}^{uu}$ be the strong unstable foliation of Φ , and $\mathcal{F}^s$ be the weak stable foliation of Φ . For each $p \in M$, we can take an open neighborhood U_p and a C^ω coordinate $\varphi_p : U_p \rightarrow \mathbb{R}^N$ such that

- $\varphi_p(p) = (0, 0) \in \mathbb{R}^{N-1} \times \mathbb{R}$,
- $\varphi_p(U_p) = (-\delta_p, \delta_p)^N$ for some $\delta_p > 0$, and
- for any $(x, y) = (x_1, \dots, x_{N-1}, y) \in \mathbb{R}^{N-1} \times \mathbb{R}$ and any sufficiently small $t \geq 0$,

$$\begin{aligned} \varphi_p^{-1}(x \times (-\delta_p, \delta_p)) &\subset \mathcal{F}^{uu}(\varphi_p^{-1}(x, 0)), \\ \varphi_p^{-1}((-\delta_p, \delta_p)^{N-1} \times y) &\subset \mathcal{F}^s(\varphi_p^{-1}(0, y)), \\ \Phi^t \circ \varphi_p^{-1}(x_1, x_2, \dots, x_{N-1}, y) &= \varphi_p^{-1}(x_1 + t, x_2, \dots, x_{N-1}, y). \end{aligned} \quad (30)$$

For each $x \in (-\delta_p, \delta_p)^{N-1}$, we define a C^ω map $l_{p,x}$ from $(-\delta_p, \delta_p)$ to $\mathcal{F}^{uu}(\varphi_p^{-1}(x, 0))$ by

$$l_{p,x}(y) = \varphi_p^{-1}(x, y).$$

Let $\{\mu_p\}_{p \in M}$ be the Margulis measure of Φ along $\mathcal{F}^{uu}$. For each $p \in M$ and $(w, x) \in (-\delta_p, \delta_p)^{N-1}$, we define a measure $\nu_{p,x}$ on $(-\delta_p, \delta_p)$ and a function ζ_p on $(-\delta_p, \delta_p)$ by

$$\begin{aligned} \nu_{p,x} &= \mu_{\varphi_p^{-1}(x, 0)} \circ l_{p,x}, \\ \zeta_p(x, y) &= \frac{d\nu_{p,x}}{d\nu_{p,0}}(y). \end{aligned}$$

By Proposition 4.2, ζ_p is real analytic for any $p \in M$.

Let $\check{y}_p$ be a function on U_p given by

$$\check{y}_p \circ \varphi_p^{-1}(x, y) = \int_0^y d\nu_{p,x} = \int_0^y \zeta_p(x, y') d\nu_{p,0}(y').$$

Since each μ_p is non-atomic and positive on each non-empty open subset of $\mathcal{F}^{uu}(p)$, we can define a *continuous* coordinate $\check{\varphi}_p : U_p \rightarrow \mathbb{R}^N$ by

$$\check{\varphi}_p(q) = (x, \check{y}_p(q)). \quad (31)$$

for any $q = \varphi_p^{-1}(x, y) \in U_p$.

By $\{Z_1, \dots, Z_N\}$, we denote the standard orthonormal frame of $\mathbb{R}^N$.

Lemma 4.4. *The family $\{\check{\varphi}_p\}_{p \in M}$ has the following properties:*

1. *The map $\check{\varphi}_p \circ \check{\varphi}_q^{-1}$ is real analytic for any $p, q \in M$.*
2. *$D(\check{\varphi}_p \circ \check{\varphi}_q^{-1})(Z_N) = Z_N$.*
3. *For any fixed $y \in (-\delta_p, \delta_p)$, the map $x \mapsto \check{\varphi}_p \circ \varphi_p(x, y)$ is also of class C^ω .*

Proof. There exist C^ω local diffeomorphisms h of $\mathbb{R}^{N-1}$ and v of $\mathbb{R}$ such that $\varphi_p \circ \varphi_q^{-1}(x, y) = (h(x), v(y))$ for any $(x, y) \in \varphi_q(U_p \cap U_q)$. To prove the first assertion, it is sufficient to show that $\check{y}_p \circ \check{\varphi}_q^{-1}$ is real analytic. Since $\mathcal{F}^{uu}(\varphi_p^{-1}(h(x), 0)) = \mathcal{F}^{uu}(\varphi_q^{-1}(x, 0))$, we have $\mu_{\varphi_p^{-1}(h(x), 0)} = \mu_{\varphi_q^{-1}(x, 0)}$. It implies $\nu_{p, h(x)} \circ v = \nu_{q, x}$. Then,

$$\begin{aligned} \check{y}_p \circ \varphi_q^{-1}(x, y') &= \int_0^{v(y')} d\nu_{p, h(x)} \\ &= \int_{v^{-1}(0)}^{y'} d\nu_{q, x} = \int_0^{y'} d\nu_{q, x} - \int_0^{v^{-1}(0)} d\nu_{q, x} \\ &= \check{y}_q \circ \varphi_q^{-1}(x, y') - \int_0^{v^{-1}(0)} d\nu_{q, x} \end{aligned}$$

for any $(x, y') \in \varphi_q(U_p \cap U_q)$. Hence,

$$\check{y}_p \circ \check{\varphi}_q^{-1}(x, y) = y - \int_0^{v^{-1}(0)} \zeta_q(x, \eta) d\nu_{q, 0}(\eta).$$

It is easy to see that $Z_N(\check{y}_p \circ \check{\varphi}_q^{-1}) = 1$ and

$$Z_{i_1} \cdots Z_{i_k}(\check{y}_p \circ \check{\varphi}_q^{-1})(x, y) = - \int_0^{v^{-1}(0)} (Z_{i_1} \cdots Z_{i_k} \zeta_q)(x, \eta) d\nu_{q, 0}(\eta). \quad (32)$$

for any $i_1, \dots, i_k = 1, \dots, N-1$. Since ζ is real analytic, also the right-hand term of the above equation is. Therefore, $\check{y}_p \circ \check{\varphi}_q^{-1}$ is real analytic.

The second assertion is an immediate consequence of the equation $Z_N(\check{y}_p \circ \check{\varphi}_q^{-1}) = 1$. The right-hand side of the equation (32) does not depend on y . It implies the last assertion. $\square$

The first assertion of the lemma implies that the family $\{\check{\varphi}_p\}_{p \in M}$ defines a C^ω structure of M . Let $\check{M}$ be the C^ω manifold such that $\check{M} = M$ as a set and its C^ω chart is given by $\{\check{\varphi}_p\}_{p \in M}$. The second assertion of the lemma implies that there exists a C^ω vector field Y^u on $\check{M}$ such that $D\check{\varphi}_p(Y^u) = Z_N$ for any $p \in M$.

Let $H : M \rightarrow \check{M}$ be the identity map as a set. We define continuous foliations $\check{\mathcal{F}}^{uu}$ and $\check{\mathcal{F}}^s$ on $\check{M}$ by $\check{\mathcal{F}}^{uu}(H(p)) = H(\mathcal{F}^{uu}(p))$ and $\check{\mathcal{F}}^s(H(p)) = H(\mathcal{F}^s(p))$. Since $\check{\varphi}_p^{-1}(0 \times \mathbb{R}) = \varphi_p^{-1}(0 \times \mathbb{R}) \subset \mathcal{F}^{uu}(p)$, the vector field Y^u is tangent to $\check{\mathcal{F}}^{uu}$. In particular, $\check{\mathcal{F}}^{uu}$ is a C^ω foliation.

Let $TM = T\Phi \oplus E^{ss} \oplus E^{uu}$ be the Anosov splitting of Φ . By the last assertion of Lemma 4.4, the restriction of H to each leaf of $\mathcal{F}^s$ is of class C^ω . Hence, the restriction of DH to $T\mathcal{F}^s = T\Phi \oplus E^{ss}$ is well-defined. Put $\check{E}^{ss} = DH(E^{ss})$ and $\check{E}^{uu} = \mathbb{R} \cdot Y^u = T\check{\mathcal{F}}^{uu}$. We define a continuous flow $\check{\Phi}$ on $\check{M}$ by $\check{\Phi}^t = H \circ \Phi^t \circ H^{-1}$. The following lemma implies that $\check{\Phi}$ is an Anosov flow satisfying the second condition in Proposition 4.3.

Lemma 4.5. *$\check{\Phi}$ is a C^ω Anosov flow such that*

- $T\check{M} = T\check{\Phi} \oplus \check{E}^{ss} \oplus \check{E}^{uu}$ is the Anosov splitting of $\check{\Phi}$, and
- $D\check{\Phi}^t(Y^u(p)) = e^{\lambda t} \cdot Y^u(\check{\Phi}^t(p))$ for any $p \in \check{M}$ and $t \in \mathbb{R}$.

Proof. By (30), we have

$$\Phi^t \circ l_{p, (x_1, x_2, \dots, x_{N-1})} = l_{p, (x_1+t, x_2, \dots, x_{N-1})}$$

for any $p \in M$, $(x_1, x_2, \dots, x_{N-1}) \in (-\delta_p, \delta_p)^{N-1}$, and $t \in \mathbb{R}$ whose absolute value is sufficiently small. Hence, the equation (24) implies

$$\check{\Phi}^t \circ \check{\varphi}_p^{-1}(x_1, x_2, \dots, x_{N-1}, y) = \check{\varphi}_p^{-1}(x_1 + t, x_2, \dots, x_{N-1}, e^{\lambda t} y). \quad (33)$$

Since $\check{\varphi}_p$ is a C^ω coordinate of $\check{M}$, the flow $\check{\Phi}$ is real analytic. Take a C^∞ metric $\|\cdot\|_1$ on $\check{M}$ such that $\|Y^u\|_1 = 1$. Since $D\check{\varphi}_p(Y^u) = Z_N$, the equation (33) implies $D\check{\Phi}^t(Y^u(p)) = e^{\lambda t} \cdot Y^u(\check{\Phi}^t(p))$. In particular, $\check{E}^{uu}$ is $D\check{\Phi}$ -invariant and $\|D\check{\Phi}^t|_{\check{E}^{uu}(p)}\|_1 = e^{\lambda t}$ for any $p \in \check{M}$ and $t \in \mathbb{R}$. Since $DH|_{E^{ss}}$ is a continuous bundle map, there exists $C > 1$ such that $C^{-1}\|v\| \leq \|Dh(v)\|_1 \leq C\|v\|$ for any $v \in E^{ss}$. It implies

$$\|D\check{\Phi}^t|_{\check{E}^{ss}(H(p))}\|_1 \leq C^2 \|D\Phi^t|_{E^{ss}(p)}\|,$$

where the right-hand term decays exponentially. Therefore, $\check{\Phi}$ is an C^ω Anosov flow with the desired properties. $\square$

Lemma 4.6. *H can be C^0 -approximated by a C^∞ diffeomorphism $\check{H}$ such that $\check{H}(\mathcal{F}^{uu}(p)) = \check{\mathcal{F}}^{uu}(H(p))$ for any $p \in M$.*

Proof. Let Ψ be a C^∞ flow on M which is generated by a unit vector field tangent to $\mathcal{F}^{uu}$. Fix a C^∞ even function g on $\mathbb{R}$ such that $g'(t) \leq 0 \leq g(t)$ for any $t \geq 0$, the support of g is compact, and the integral of g over $\mathbb{R}$ is one. For $\epsilon > 0$ and a given continuous function f on M , we define a function $\mathcal{S}_\epsilon f$ on M by

$$(\mathcal{S}_\epsilon f)(p) = \int_{\mathbb{R}} (f \circ \Psi^{\epsilon\tau})(p) \cdot g(\tau) d\tau.$$

Take a finite subset $\{p_1, \dots, p_m\}$ of M so that $M = \bigcup_{i=1}^m U_{p_i}$. Let $\{\chi_i\}_{i=1}^m$ be a partition of unity associated with the covering $\{U_{p_i}\}_{i=1}^m$ of M . For each $i = 1, \dots, m$, we define a map π_i from U_{p_i} to $\mathcal{F}^s(p_i)$ by

$$\pi_i \circ \varphi_{p_i}^{-1}(x, y) = \varphi_{p_i}^{-1}(x, 0).$$

We assume the following claim, whose proof is postponed until we finish the proof of the lemma.

Claim 4.7. $\mathcal{S}_\epsilon(\chi_i \cdot \check{y}_{p_i})$ is a C^∞ function on M .

Now, the proof of the lemma is almost identical to Lemma in [21] by de la Llave and Moriyón. Let $\check{\Psi}$ be the C^∞ flow on $\check{M}$ which is generated by Y^u . Since $D\check{\varphi}_p(Y^u) = Z_N$ and $\check{y}_p \circ \check{\varphi}_p^{-1}(x, y) = y$, we have

$$p = \check{\Psi}^{\check{y}_{p_i}(p)}(H \circ \pi_i(p)) \quad (34)$$

for any $p \in U_{p_i}$. For each small $\epsilon > 0$, we define a map H_ϵ from M to $\check{M}$ by

$$H_\epsilon(p) = \check{\Psi}^{(\mathcal{S}_\epsilon - I)(\sum_{i=1}^m \chi_i \cdot \check{y}_{p_i})(p)}(H(p)),$$

where I is the identity operator. Remark that $H_\epsilon(\mathcal{F}^{uu}(p)) = \check{\mathcal{F}}^{uu}(H(p))$ for any $p \in M$.

For any $p \in U_{p_i} \cap U_{p_j}$, the equation (34) implies

$$\check{y}_{p_i}(p) = \check{y}_{p_j}(p) + \check{y}_{p_i} \circ \pi_j(p).$$

Hence, we have

$$\begin{aligned} \check{\Psi}^{-\sum_{i=1}^m \chi_i \cdot \check{y}_{p_i}(p)}(H(p)) &= \check{\Psi}^{-\sum_{i=1}^m \chi_i \cdot (\check{y}_{p_i} \circ \pi_j)(p) - \check{y}_{p_j}(p)}(H(p)) \\ &= \check{\Psi}^{-\sum_{i=1}^m \chi_i \cdot (\check{y}_{p_i} \circ \pi_j)(p)}(H \circ \pi_j(p)) \end{aligned}$$

for any $p \in U_{p_j}$. Since π_j is a C^∞ map from U_{p_j} to $\mathcal{F}^s(p_j)$ and the restrictions of $\check{y}_{p_i}$ and H to a leaf of $\mathcal{F}^s$ are of class C^∞ , the last term depends C^∞ -differentiably with respect to p . By Claim 4.7,

$$H_\epsilon(p) = \check{\Psi}^{\mathcal{S}_\epsilon(\sum_{i=1}^m \chi_i \cdot \check{y}_{p_i})(p)} \circ \check{\Psi}^{-\sum_{i=1}^m \chi_i \cdot (\check{y}_{p_i} \circ \pi_j)(p)}(H \circ \pi_j(p)).$$

is smooth with respect to p . By the same argument as [21], we can see that H_ϵ is a diffeomorphism and is C^0 -close to H if ϵ is sufficiently small. $\square$

Proof of Claim 4.7. Fix $i = 1, \dots, m$ and $\epsilon > 0$. Put $f = \chi \cdot \check{y}_{p_i}$. We will show that $(\mathcal{S}_\epsilon f) \circ \varphi_{p_i}^{-1}$ is a C^∞ function. By Journé's theorem [17], it is sufficient to show that $Z_{i_1} \cdots Z_{i_k}(\mathcal{S}_\epsilon f) \circ \varphi_{p_i}^{-1}$ and $Z_N^k(\mathcal{S}_\epsilon f) \circ \varphi_{p_i}^{-1}$ are Hölder continuous for any $k \geq 1$ and any $i_1, \dots, i_k = 1, \dots, N-1$.

Let U be a neighborhood of the support of $f \circ \varphi_{p_i}^{-1}$ whose closure is contained in $\varphi_{p_i}(U_p)$. There exists a small neighborhood I of 0 in $\mathbb{R}$ and a smooth function $\bar{\tau}$ on $U \times I$ such that

$$\varphi_{p_i}^{-1}(x, y + \eta) = \Psi^{\epsilon \cdot \bar{\tau}(x, y, \eta)}(\varphi_{p_i}^{-1}(x, y))$$

for any $(x, y) \in U \subset \mathbb{R}^{N-1} \times \mathbb{R}$ and $\eta \in I$. Then, we have

$$\begin{aligned} (\mathcal{S}_\epsilon f) \circ \varphi_{p_i}^{-1}(x, y) &= \int_{\mathbb{R}} (f \circ \Psi^{\epsilon \tau} \circ \varphi_{p_i}^{-1})(x, y) \cdot g(\tau) d\tau \\ &= \int_{\mathbb{R}} (f \circ \varphi_{p_i}^{-1})(x, y + \eta) \cdot \left((g \circ \bar{\tau}) \cdot \frac{\partial \bar{\tau}}{\partial \eta} \right)(x, y, \eta) d\eta. \end{aligned}$$

It is known that any topological conjugacy between two Anosov flows is a bi-Hölder homeomorphism (see *e.g.* Theorem 19.1.2 of [18]). Hence, the map $\tilde{\varphi}_{p_i} \circ \varphi_{p_i}^{-1}$ is Hölder continuous. By (32), also $Z_{i_1} \cdots Z_{i_k}(\tilde{y}_{p_i} \circ \varphi_{p_i}^{-1})$ is for any $i_1, \dots, i_k = 1, \dots, N-1$. Since χ, g and $\bar{\tau}$ are of class C^∞ , the above equation implies that $Z_{i_1} \cdots Z_{i_k}(\mathcal{S}_\epsilon f \circ \varphi_{p_i}^{-1})$ is Hölder continuous for any $i_1, \dots, i_k = 1, \dots, N_1$. By an easy calculation, we can see that $Z_N^k(\mathcal{S}_\epsilon f \circ \varphi_{p_i}^{-1})$ is also a Hölder continuous function for any $k \geq 1$. As mentioned at the beginning of the proof, Journé's theorem [17] implies that $\mathcal{S}_\epsilon f$ is of class C^∞ . $\square$

Now, we prove Proposition 4.3. The equation (28) follows from Lemma 4.5. Let $\tilde{H}$ be the diffeomorphism in Lemma 4.6. Since $\tilde{H}(p) \in \tilde{\mathcal{F}}^{uu}(H(p)) \cap \tilde{H}(\mathcal{F}^{uu}(p))$, we have $\tilde{H}(\mathcal{F}^{uu}(p)) = \tilde{\mathcal{F}}^{uu}(H(p)) = \tilde{\mathcal{F}}^{uu}(\tilde{H}(p))$. It implies that

$$\begin{aligned} \tilde{H} \circ \Phi^t(\mathcal{F}^{uu}(p)) &= \tilde{H}(\mathcal{F}^{uu}(\Phi^t(p))) = \tilde{\mathcal{F}}^{uu}(H \circ \Phi^t(p)) = \tilde{\mathcal{F}}^{uu}(\tilde{\Phi}^t \circ H(p)) \\ &= \tilde{\Phi}^t(\tilde{\mathcal{F}}^{uu}(H(p))) = \tilde{\Phi}^t(\tilde{\mathcal{F}}^{uu}(\tilde{H}(p))). \end{aligned}$$

Therefore, the equation (27) holds.

4.3 Proof of Theorems B and C

Now, we show Theorems B and C. By Propositions 3.10, it is sufficient to construct an action ρ in $\mathcal{A}_\Gamma$ with $\bar{a}(\rho) = a$ for any give $a \in \Delta_\Gamma$ and show it is C^∞ conjugate to a C^ω action.

Fix $a \in \Delta_\Gamma$. Let Φ_Γ be the homogeneous Anosov flow on M_Γ and $\mathcal{F}_\Gamma^u$ be its weak unstable foliation. By Propositions 3.5 and 3.6, there exists a C^ω Anosov flow Φ_a on M_Γ such that its Anosov splitting is real analytic, the orbits of Φ_a coincide with those of Φ_Γ , and

$$\tau(p; \Phi_a) = \tau(p; \Phi_\Gamma) + a(\gamma_\Gamma(p)) \quad (35)$$

for any $p \in \text{Per}(\Phi)$. Remark that the weak unstable foliations of Φ_a is $\mathcal{F}_\Gamma^u$ since Φ_a and Φ_Γ have the same orbits.

Let $\tilde{\Phi}, \tilde{M}, Y^u, \tilde{H}$, and λ be the objects obtained by applying Proposition 4.3 to Φ_a . The vector field Y^u generates a C^ω flow $\tilde{\Psi}$ on $\tilde{M}$. The equation (28) implies the identity $\tilde{\Phi}^t \circ \tilde{\Psi}^y = \tilde{\Psi}^{e^{\lambda t} y} \circ \tilde{\Phi}^t$. We define actions $\check{\rho} : \tilde{M} \times \text{GA} \rightarrow \tilde{M}$ and $\rho : M \times \text{GA} \rightarrow M$ by

$$\begin{aligned} \check{\rho}(p, X^t U^y) &= \tilde{\Psi}^y \circ \tilde{\Phi}^{\frac{t}{\lambda}}(p), \\ \rho(p, X^t U^y) &= \tilde{H}^{-1} \circ \check{\rho}(\tilde{H}(p), X^t U^y). \end{aligned}$$

The action ρ on M is real analytic. By the equation (27), the orbit foliation of ρ coincides with the weak unstable foliation of Φ_a . In particular, ρ is an element of $\mathcal{A}_\Gamma$ and $\Phi_\rho^t = \tilde{H}^{-1} \circ \tilde{\Phi}^{\frac{t}{\lambda}} \circ \tilde{H}$.

The equation (29) implies $\tau(H_\rho(p); \Phi_\rho) = \lambda \cdot \tau(p; \Phi_a)$ for any $p \in \text{Per}(\Phi_\Gamma)$. By the equations (16) and (35), we have

$$J(H_\rho(p); \Phi_\rho) = (1 - \lambda^{-1})\tau(H_\rho(p); \Phi_\rho) + a(\gamma_\Gamma(p)). \quad (36)$$

By taking $p \in \text{Per}(\Phi_\Gamma)$ whose orbit is null-homologous, we obtain $\bar{\delta}(\rho) = 1 - \lambda^{-1}$. Since $H^1(M_\Gamma, \mathbb{R})$ is spanned by periodic orbits of Φ_Γ , the above equation implies that $\bar{a}(\rho) = a$.

5 Proof of Theorem D

We recall a classification of Anosov flows with the C^2 Anosov splitting due to Ghys [13, 14].

Theorem 5.1 (Ghys [13, 14]). *Let Γ be a cocompact lattice of $\widetilde{\text{PSL}}(2, \mathbb{R})$ and Φ be a C^∞ Anosov flow on M_Γ . If Φ admits a C^2 Anosov splitting, then there exist cocompact lattices Γ_s and Γ_u which are isomorphic to Γ and such that the weak stable foliation of Φ is C^∞ diffeomorphic to $\mathcal{F}_{\Gamma_s}^s$ and the weak unstable foliation of Φ is C^∞ diffeomorphic to $\mathcal{F}_{\Gamma_s}^u$.*

Now, we prove Theorem D. Let ρ be an action in $\mathcal{A}_\Gamma$ for some cocompact lattice Γ of $\widetilde{\text{PSL}}(2, \mathbb{R})$. Put $a = \bar{a}(\rho)$ and $\delta = \bar{\delta}(\rho)$. By Lemma 2.10, for any given $\gamma \in \Gamma$ with $L(\gamma) \neq 0$, there exist $p_\gamma \in \text{Per}(\Phi_\Gamma)$, $c \in \Gamma$, and $n_\gamma \geq 1$ such that c is central and $\gamma_\Gamma(p_\gamma) = c \cdot \gamma^{n_\gamma}$.

Suppose that the weak stable foliation $\mathcal{F}^s$ of Φ is of class C^2 . By the above theorem, there exists a cocompact lattice Γ' and a C^2 -diffeomorphism $H : M_\Gamma \rightarrow M_{\Gamma'}$ such that $H(\mathcal{F}^s) = \mathcal{F}_{\Gamma'}^s$. It induces an isomorphism $\sigma : \Gamma \rightarrow \Gamma'$ between the fundamental groups of M_Γ and $M_{\Gamma'}$.

Let $\Phi_{\Gamma'}$ and Φ_1 be flows on $M_{\Gamma'}$ defined by $\Phi_{\Gamma'}^t(p) = \rho_{\Gamma'}(p, X^t)$ and $\Phi_1^t = H \circ \Phi^t \circ H^{-1}$. They have the same weak stable foliations $\mathcal{F}_{\Gamma'}^s$. By Proposition 3.4, there exists a topological equivalence H' between $\Phi_{\Gamma'}$ and Φ_1 which is isotopic to the identity and preserves each leaf of $\mathcal{F}_{\Gamma'}^s$. Then, $H_{\Gamma, \Gamma'} = H' \circ H \circ H_\rho$ is a topological equivalence between Φ_Γ and $\Phi_{\Gamma'}$. It satisfies $\gamma_{\Gamma'}(H_{\Gamma, \Gamma'}(p_\gamma)) = \sigma(\gamma_\Gamma(p_\gamma))$, and hence,

$$J^u(H_\rho(p_\gamma); \Phi_\rho) = J^u(H \circ H_\rho(p_\gamma); \Phi_1) = J^u(H_{\Gamma, \Gamma'}(p_\gamma); \Phi_{\Gamma'}) = L(\sigma(\gamma_\Gamma(p_\gamma))).$$

By the equations (14), (15), and (19), we have

$$\begin{aligned} n_\gamma \cdot a(\gamma) &= a(\gamma_\Gamma(p_\gamma)) = (1 - \delta)\tau(H_\rho(p_\gamma), \Phi_\rho) - \tau(p_\gamma, \Phi_\Gamma) \\ &= (1 - \delta)L(\sigma(\gamma_\Gamma(p_\gamma))) - L(\gamma_\Gamma(p_\gamma)) \\ &= n_\gamma \{(1 - \delta)L(\sigma(\gamma)) - L(\gamma)\}. \end{aligned} \tag{37}$$

Fix a hyperbolic element $\gamma \in \Gamma$. Since Γ is cocompact and $\gamma_\Gamma(p_\gamma)$ is hyperbolic, there exists $\gamma' \in \Gamma$ such that $\eta_m = [(\gamma_\Gamma(p_\gamma))^m, \gamma']$ is not contained in the center for any $m \geq 1$. By Proposition 2.7, η_m is not parabolic. The equation 37 implies

$$(1 - \delta)L(\sigma(\eta_m)) - L(\eta_m) = 0.$$

for any $m \geq 1$. By Lemma 2.8, we have $(1 - \delta)L(\sigma(\gamma_\Gamma(p_\gamma))) = L(\gamma_\Gamma(p_\gamma))$, and hence, $a(\gamma) = 0$ by (37). Therefore, $a = \bar{a}(\rho)$ is the zero class. By Theorem A, ρ is homogeneous.

A Locally free actions on solvable manifolds

In this appendix, we give a proof of the following unpublished result.

Theorem A.1 (Ghys). *Let M be a closed three-dimensional manifold whose fundamental group is solvable. Then, any C^∞ locally free action of GA on M preserves a continuous volume. In particular, it is homogeneous.*

Proof. Let ρ be a C^∞ locally free action of GA on M . We define a flow Φ on M by $\Phi^t(p) = \rho(p, X^t)$. By Theorem 1.5, it is Anosov. Let X be the vector field generating Φ and $TM = T\Phi \oplus E^{ss} \oplus E^{uu}$ be the Anosov splitting of Φ . Plante [24] showed that any Anosov flow on a solvable three-dimensional manifolds is topologically equivalent to the suspension of a hyperbolic toral automorphism. Hence, $H_1(M, \mathbb{R})$ is one-dimensional, and there exists a smooth closed one-form ω on M such that $\omega(X)$ is positive everywhere and ω represents a generator of $H^1(M, \mathbb{R})$.

For each $p \in \text{Per}(\Phi)$ we denote the homology class represented by the orbit of p by c_p . By a classification of smooth foliations on the mapping torus of the hyperbolic toral automorphism [15], the orbit foliation of ρ is diffeomorphic to the suspension of a linear foliation of the torus. Hence, there exists $\lambda^s < 0$ such that $J^s(p; \Phi) = \lambda^s \cdot \omega(c_p)$ for any $p \in \text{Per}(\Phi)$. It implies

$$J(p; \Phi) = J^u(p; \Phi) + J^s(p; \Phi) = \tau(p; \Phi) + \lambda^s \omega(c_p). \quad (38)$$

Take $\lambda \in \mathbb{R}$ so that $\lambda \cdot \omega$ represents the class $\bar{a}(\rho)$. The same argument as in Subsection 3.3 implies that there exists a constant $\delta < 1$ such that

$$J(p; \Phi) = \delta \cdot \tau(p; \Phi) + \lambda \cdot \omega(c_p) \quad (39)$$

for any $p \in \text{Per}(\Phi)$. Put $\lambda_* = (\lambda - \delta \lambda^s)(1 - \delta)^{-1}$. By the equations (38) and (39), we have $J(p; \Phi) = \lambda_* \cdot \omega(c_p)$ for any $p \in \text{Per}(\Phi)$. The C^∞ -Livschitz theorem implies that there exists a smooth volume v on TM such that

$$\log \det_v D\Phi^t(p) = \lambda_* \int_0^t \omega(X) \circ \Phi^s(p) ds$$

for any $P \in M$ and $t \in \mathbb{R}$, where $\det_v$ is the determinant with respect to the volume v . By integrating the exponential of both sides on M , we obtain that

$$1 = \frac{v(\Phi^t(M))}{v(M)} = \int_M \exp \left(\lambda_* \int_0^t \omega(X) \circ \Phi^s(p) ds \right) dv(p).$$

Since $\omega(X)$ is positive everywhere, it implies that $\lambda_* = 0$. Therefore, Φ preserves the volume v . $\square$

References

- [1] M. Asaoka, Regular projectively Anosov flows on three-dimensional manifolds. preprint (arXiv:math.DS/0504208).

- [2] A.F. Beardon, The geometry of discrete groups. Graduate Texts in Mathematics, 91. *Springer-Verlag*, New York, 1995.
- [3] M. Belliard, Actions localement libres de groupes non unimodulaires. *Ann. Fac. Sci. Toulouse Math. (6)* **7** (1998), no. 1, 35–50.
- [4] M. Belliard, Locally free actions of solvable Lie groups and cohomology. *Ergodic Theory Dynam. Systems* **26** (2006), no. 1, 1–17.
- [5] M. Belliard, A differentiable classification of certain locally free actions of solvable Lie groups. preprint.
- [6] M. Belliard and O. Birembaux, Actions localement libres de groupes résolubles. *Ann. Inst. Fourier (Grenoble)* **44** (1994), no. 5, 1519–1537.
- [7] E. E. Cawley, The Teichmüller space of an Anosov diffeomorphism of T^2 . *Invent. Math.* **112** (1993), no. 2, 351–376.
- [8] N. Chernov, Invariant measures for hyperbolic dynamical systems. *Handbook of dynamical systems, Vol. 1A*, 321–407, North-Holland, Amsterdam, 2002.
- [9] É. Ghys, Sur les actions localement libres du group affine. Thèse de 3ème cycle, Lille 1979.
- [10] É. Ghys, Flots d’Anosov sur les 3-variétés fibrées en cercles. *Ergodic Theory Dynam. Systems* **4** (1984), no. 1, 67–80.
- [11] É. Ghys, Actions localement libres du groupe affine. *Invent. Math.* **82** (1985), no. 3, 479–526.
- [12] É. Ghys, Flots d’Anosov dont les feuilletages stables sont différentiables. *Ann. scient. Éc. Norm. Sup. (4)* **20** (1987), no. 2, 251–270.
- [13] É. Ghys, Rigidité différentiable des groupes fuchsien. *Inst. Hautes Études Sci. Publ. Math.* **78** (1993), 163–185.
- [14] É. Ghys, Déformations de flots d’Anosov et de groupes fuchsien. *Ann. Inst. Fourier (Grenoble)* **42** (1992), no. 1–2, 209–247.
- [15] É. Ghys and V. Sergiescu. Stabilité et conjugaison différentiable pour certains feuilletages. *Topology* **19** (1980), no. 2, 179–197.
- [16] B. Hasselblatt and A. Katok, Principal structures. *Handbook of dynamical systems, Vol. 1A*, 1–203, North-Holland, Amsterdam, 2002.
- [17] J.L. Journé, On a regularity problem occurring in connection with Anosov diffeomorphisms. *Commun. Math. Phys.* **106** (1986), no.2, 345–351.
- [18] A. Katok and B. Hasselblatt, Introduction to the modern theory of dynamical systems. Encyclopedia of Mathematics and its Applications, 54. *Cambridge University Press, Cambridge*, 1995.

- [19] Livšic, A. N., Certain properties of the homology of Y -systems. *Math. Notes* **10** (1971), 758–763. (translated from *Mat. Zametki* **10** (1971), 555–564)
- [20] R. de la Llave, J.M. Marco, and R.Moriyón, Canonical perturbation theory of Anosov systems and regularity results for the Livšic cohomology equation. *Ann. of Math. (2)* **123** (1986), no. 3, 537–611.
- [21] R. de la Llave and R. Moriyón, Invariants for smooth conjugacy of hyperbolic dynamical systems. IV. . *Commun. Math. Phys.* **116** (1988), no.2, 185–192.
- [22] G. A. Margulis, On some aspects of the theory of Anosov systems. With a survey by Richard Sharp: Periodic orbits of hyperbolic flows. Translated from the Russian by Valentina Vladimirovna Szulikowska. Springer Monographs in Mathematics. *Springer-Verlag, Berlin*, 2004.
- [23] Y. Mitsumatsu and S. Matsumoto, Leafwise cohomology and rigidity of certain Lie group actions. *Ergodic Theory Dynam. Systems* **23** (2003), no. 6, 1839–1866.
- [24] J.F. Plante, Anosov flows, transversely affine foliations, and a conjecture of Verjovsky. *J. London Math. Soc. (2)* **23** (1981), no. 2, 359–362.
- [25] F. Salein, Variétés anti-de Sitter de dimension 3 possédant un champ de Killing non trivial. *C. R. Acad. Sci. Paris Sér. I Math.* **324** (1997), no. 5, 525–530.
- [26] K. Sigmund, Generic properties of invariant measures for Axiom A diffeomorphisms. *Invent. Math.* **11** (1970), 99–109.
- [27] N. Tsuchiya and A. Yamakawa, Rigidity of codimension one locally free actions of solvable Lie groups. *Current trends in transformation groups*, 181–200, *K-Monogr. Math.*, 7, *Kluwer Acad. Publ., Dordrecht*, 2002.
- [28] A. Yamakawa and N. Tsuchiya, Codimension one locally free actions of solvable Lie groups. *Tohoku Math. J. (2)* **53** (2001), no. 2, 241–263.